

Citrona™ Series Erysimum

These sweet-smelling “Pansy Pals” provide a low-maintenance option for Autumn programs!

- **Citrona™** Erysimum does not require vernalization and can be grown in tandem with pansies in cooler temperatures.
- Grows in just 11 to 12 weeks with strong branching, making these plants perfect for large packs and small pot programs.
- **Citrona™** Erysimum can be scheduled for multiple programs throughout the year.
- Contact your local distributor or call PanAmerican Seed at 800 231-4868 for ordering information on **Citrona™** Erysimum.

Erysimum allionii

Approximate seed count: 14,000 S./oz. (500 S./g.)

Plug Production

Plug Tray Size

Can be produced in 392 or 406 cell-type plug trays. The average plug finishing time is 4 weeks.

Media

Use a well-drained, disease-free plug media with a pH of 5.8 to 6.2.

Sowing

Cover the seed with a light to medium layer of coarse grade vermiculite. Because of **Citrona's** excellent basal branching, these plants can be sown single seeded in contrast to others in the marketplace.

Days to Germ: 2 to 4 days.

Stage 2 timing: 7 days

Stage 3 timing: 10 days

Stage 4 timing: 7 days

Temperature

Germination: 68 to 72°F (20 to 22°C).

Stage 2: 65 to 70°F (18 to 21°C) days; 60°F (15°C) nights

Stage 3: 65°F (18°C) days; 60°F (15°C) nights

Stage 4: 60°F (15°C) days; 55°F (13°C) nights

Light

Light is not required for germination. For growing on the plugs, light levels can be up to 3,000 f.c. (30,000 Lux).

Humidity

Maintain 95 to 97% relative humidity during germination.

Fertilizer

Beginning at Stage 3, start fertilizing the seedlings twice a week with 50 ppm N, using a complete balanced plug fertilizer. Increase the nitrogen concentration to 100 ppm after a week, and continue this regime until finishing the plugs. Maintain an EC of 0.5 to 0.75 mmhos/cm and a pH of 5.8 to 6.2 at Stage 2; at Stage 3 and 4, EC and pH values can be at 1.0 and 5.8 to 6.2, respectively.

Growth Regulators

Plant Growth Regulators are recommended for plug production. Use foliar sprays of Bonzi at 10 ppm at Week 3 from sowing to produce compact/stretch-free seedlings. One application will be sufficient during the plug production.

Growing On to Finish

Container Size

306 Premium Packs, 4 in. (10 cm), 4.5 in. (11 cm) and larger containers.

Media

Use a well-drained, disease-free soilless growing media with a pH of 5.8 to 6.2 and a medium level of initial nutrient charge.

Temperature

Erysimum performs well when grown under cooler temperatures, more like pansies, and can also be a good Autumn companion for pansies. After transplant, the crop can be grown under greenhouse conditions (65 to 70°F/18 to 21°C days and 55 to 60°F/13 to 16°C nights) for approximately 2 weeks until the plugs start rooting out and are well established, after which they can be grown outside under cold frame-type conditions to flower or finish. The optimal outside growing temperatures can be 65 to 75°F (18 to 29°C) days and 40 to 55°F (5 to 13°C) nights.

Light

No supplemental lighting is required.

Irrigation & Fertilization

Maintain even moisture and avoid excessive wetness. Fertilize the crop weekly with 200 to 300 ppm N with a complete balanced fertilizer, maintaining a pH of 5.8 to 6.2. E.C. of the media can be at 1.5 to 2.5 mmhos/cm (SME).

Plant Growth Regulators

Growth Regulators can be used for pack/container production. Foliar sprays of Bonzi at 20 ppm can be applied 1 to 2 times after transplant. First application can be done 10 to 14 days after transplant, followed by a second application 2 weeks later if necessary (especially if the crop is targeted for finishing in Autumn). Alternatively, a tank mix foliar sprays of B-Nine at 2,500 ppm and Cycocel at 1,000 ppm can also be effective. The timing and frequency for the tank mix regime can be same as that of Bonzi.

Crop Scheduling

Sow to transplant: 4 weeks in a 392 or 406 plug tray

Transplant to finish in 306 cell pack: 7 to 9 weeks seasonally

Total crop time to flower from sowing: 11 to 13 weeks seasonally

Note: Orange typically flowers slightly earlier than yellow.

Garden Height

18 to 24 in. (46 to 61 cm)

Garden Spread

12 to 18 in. (30 to 46 cm)

Garden Location

Full sun

Garden Spacing (Planting Density)

8 in. (20 cm)

Common Problems

Diseases: Having a well drained, disease-free/sterilized soil and good plant spacing is important for avoiding disease in the landscapes.

PanAmerican Seed™

PanAmSeed.com

PanAmerican Seed Co.
622 Town Road
West Chicago, Illinois USA 60185-2698
630 231-1400
Fax: 630 231-3609

PanAmerican Seed Europe BV
Lavendelweg 10
NL-1435 EW Rijsenhout, Holland
+31 (0)297-383038
Fax: +31 (0)297-383036