



Ballerina Armeria

A. pseudarmeria

Approximate seed count (raw): 28,350 to 31,200 S./oz. (1,000 to 1,100 S./g)

Key flowering facts:

- First year-flowering perennial
- Photoperiod response: day-neutral plant
- Vernalization: not required
- Flowering time: early Spring

Plug Production

Media

Use a well-drained, disease-free, soilless media with a pH of 5.8 to 6.2 and a medium initial nutrient charge (EC 0.75 mmhos/cm).

Sowing

Sow 2 to 4 seeds per cell in 288 or larger plug trays. Do not cover the seeds.

Stage 1 – Germination takes 3 to 6 days

Soil temperature: 60 to 65°F (16 to 18°C)

Light: Light is optional.

Moisture: Keep soil wet (level 4) during Stage 1.

Humidity: Maintain 95 to 97%+ relative humidity (RH) until radicles emerge.

Stage 2

Soil temperature: 60 to 65°F (16 to 18°C)

Light: Can be up to 2,500 f.c. (26,900 Lux).

Moisture: Reduce soil moisture slightly (level 3 to 4) to allow the roots to penetrate into the media.

Fertilizer: Apply fertilizer at rate 1 (less than 100 ppm N; less than 0.7 mmhos/cm EC) from nitrate-form fertilizers with low phosphorous.

Stage 3

Soil temperature: 60 to 65°F (16 to 18°C)

Light: Can be up to 2,500 f.c. (26,900 Lux).

Moisture: Allow the soil moisture to dry to level 3.

Fertilizer: Increase fertilizer to rate 2 (100 to 175 ppm N; 0.7 to 1.2 mmhos/cm EC).

Growth Regulators: Generally not needed

Stage 4

Soil temperature: 60 to 65°F (16 to 18°C)

Light: Can be up to 5,000 f.c. (54,000 Lux).

Moisture: Same as Stage 3.

Fertilizer: Same as Stage 3.

Growing On to Finish

Container Size

4 to 5-in. (10 to 13-cm) square/quart pots: 1 plug per pot

Gallon (7 in./18 cm): 3 plugs per pot

Media

Use a well-drained, disease-free, soilless media with a pH of 5.8 to 6.2 and a medium initial nutrient charge (EC 0.75 mmhos/cm).

Temperature (optimum)

Nights: 50 to 58°F (10 to 14°C)

Days: 60 to 65°F (16 to 18°C)

NOTE: For early Spring sales, grow in a frost-free greenhouse/tunnel.

Light

No additional light is required.

Photoperiod

Ballerina is a day-neutral plant.

Irrigation

Keep media moisture dry (level 2; substrate color is light brown) to medium moist (level 3; substrate color is brown to dark brown). Ballerina is relatively drought tolerant.

For overwinter production, grow plants on the dry side during cold period as overwatering could result in plant loss from root rot.

Fertilizer

Ballerina generally needs low to medium fertilization. Apply fertilizer at rate 1 to 2 (70 to 175 ppm N; 0.5 to 1.2 mmhos/cm)

After overwintering, apply an extra fertilizer application (150 to 175 ppm Nitrate fertilizer; 1.0 to 1.2 mmhos/cm) when plants start to grow in early Spring.

Growth Regulators

Not needed.

Pinching

Not needed.

Crop Scheduling

Sow to transplant (288 cell plug): 5 to 6 weeks

Annual

Transplant to flower: 12 to 15 weeks

Total crop time: 16 to 21 weeks

Overwintered frost-free

Transplant to flower: 28 to 36 weeks

Total crop time: 32 to 40 weeks

Spring production:

Sow January to mid-March for natural flowering from mid-May to July.

Overwinter production: Sow late July to late August for natural flowering late May to early June of the following year.

Note: Plants from overwinter production will have a better plant habit with any more flowering stems per plant (approximately 20 to 30 stems per plant) .

Common Problems

Insect: Sciara in plug stage

Disease: Colletotrichum

Physiological: When grown too cold during growing season, leaves may turn red-purplish and could twist.

Garden and Landscape Information

- Armeria is first year-flowering perennial to USDA Hardiness Zones 7 to 9.
- Low maintenance and salt tolerant

Note: Growers should use the information presented here as a starting point. Crop times will vary depending on the climate, location, time of year, and greenhouse environmental conditions. Chemical and PGR recommendations are only guidelines. It is the responsibility of the applicator to read and follow all the current label directions for the specific chemical being used in accordance with all regulations.

United States: 630 231-1400
Europe: +31 (0)228 54 1844
kiefert-pro-seeds.com

