

Primlet™ Series Primula acaulis

Eye-Catching Mini-Rosebud Blooms in Bright & One-of-a-Kind Colors...Novelty Sells!

- Mid-season *P. acaulis* makes a great gift plant or use it in mixed combo planters
- Wide range of strong and unique colors are great for late Winter and Spring sales
- Long shelf life – up to 3 weeks – for extended sales opportunities and satisfied consumers
- Sweet fragrance
- No special culture needed – grow just like a standard *P. acaulis* type
- Contact your local distributor or call PanAmerican Seed at 800 231-4868 for ordering information on **Primlet** primulas.

P. acaulis

Approximate Seed Count (raw): 28,000/oz. (1,300/g)

Plug Production

Stage 1

- Germination: 7 to 10 days at 64°F (17°C)
- Recommended plug sizes are 512 to 72
- Cover seed lightly with coarse vermiculite to maintain moisture levels
- 10 f.c. (100 Lux) of light benefits germination, but is not required
- Provide high moisture but do not saturate (4+)

Stage 2

- Maintain light levels below 1,500 f.c. (15,000 Lux)
- Begin fertilizing with 14-0-14 at 100 ppm
- Maintain medium moisture (3 to 4)
- Maintain temperatures at 64 to 68°F (17 to 20°C)

Stage 3

- Increase light to 2,000 to 2,500 f.c. (20,000 to 25,000 Lux)
- Increase fertilizer to 200 ppm, alternating 14-0-14 and 20-10-20
- Allow moisture levels to alternate between 1 (dry, not wilted) to 4 (moist)
- Keep media pH below 6.0
- Maintain temperatures at 64 to 68°F (17 to 20°C)

Stage 4

- Increase light to 2,500 f.c. (25,000 Lux), maintaining cool temperatures
- Maintain dry to medium moisture levels

- If pH is above 6.0, apply Iron Sulfate at 1 pound per 100 gallons as a soil drench
- Maintain temperatures at 64 to 68°F (17 to 20°C)

Plug Crop Times

512/406-cell sizes: 5 to 6 weeks

288-cell size: 6 to 7 weeks

Important Plug Production Tips

- Maintain high light levels, but shade if necessary to avoid high temperatures
- Severe wilting can cause leaf tip burn
- Primula have a high pH sensitivity. Fe deficiency causes interveinal chlorosis on new growth and stunting. Maintain pH below 6.0.
- Watch for fungus gnats, especially if staying wet due to cool temperatures or shading
- Apply preventative fungicide for *Pythium* and *Theileviopsis* during late Stage 3 and early Stage 4

Growing On to Finish

Media

Use a peat-lite mix with good drainage; peat should be 50 to 60% of the mix. Maintain pH between 5.5 and 6.2. Plants in pots should be placed on a surface that allows good drainage.

Planting

Plant in moist growing media. Set the plant even or slightly above the soil line of the liner – *do not bury the plug or liner.*

Temperature

Weeks 1 to 4 after transplant (Establishing stage – goal is to establish 10 leaves)

- Nights: 55 to 60°F (12 to 15.5°C)
- Days: 60 to 65°F (15.5 to 18°C)

Weeks 5 to 10 (Bud initiation stage)

- Nights: 35 to 45°F (2 to 7°C)
- Days: 45 to 48°F (7 to 9°C)

Weeks 11 to 16 (Flower development and forcing)

- Nights: 56 to 58°F (13 to 14.5°C)
- Days: 60 to 62°F (15.5 to 16.5°C)

NOTE: Plants can be held at 40 to 45°F (4.5 to 7°C) for later forcing.

Light

Provide full sun at lower temperatures. Reduce light levels to avoid high temperatures. Excessive high light may lead to 'blind plants' or bud abortion. Optimum light is usually between 2,000 and 3,500 f.c. (20,000 and 35,000 Lux).

Humidity

Relative humidity between 60 and 70% prevents stress on the plant and reduces the water requirement. Provide good air circulation in the area to prevent condensation and *Botrytis*.

Water

Water quality should be good with alkalinity below 140 ppm. Maintain EC below 0.5 mmhos.

Fertilization

Primula is a cool crop and does not have a high fertilizer requirement. Start the plants with 20-10-20 at 200 ppm. When ready for cold treatment, change fertilizer to 15-0-15 at 50 ppm. Maintain media EC under 1.2 mmhos.

Crop Scheduling

Plug timing (based on 288-plug size): 6 to 7 weeks

Plant establishment: 4 weeks

Bud initiation and cooling: 6 weeks

Flower development and forcing: 5 to 6 weeks

Total crop time from sow to flower: 22 weeks

NOTE: Growing-on time in weeks depends on how large a plant is required. A large plant requires a longer time at 60 to 65°F (15.5 to 18°C) nights. From bud visibility to first opening of flower is approximately 4 to 5 weeks, depending on temperature.

Common Problems

Insects: Aphids, whitefly, fungus gnats

Diseases: *Botrytis* on flowers, *Pythium*, *Rhizoctonia*

Common Problem	Causes
<i>Botrytis</i>	<ul style="list-style-type: none">Plants have wet foliage and flowers at nightLack of circulating air which can prevent condensationNo fungicide control for <i>Botrytis</i>Be sure to water early in morning and avoid high humidity
Premature bud set and small plants	<ul style="list-style-type: none">Temperature too cold during growing-on periodLow fertilityNot enough weeks of warm growing-on temperatures
Flower stems too short	<ul style="list-style-type: none">Cold temperature below 40°F (4.5°C) for too longForcing at high temperatures above 65°F (18°C) nights
Flower stems long and weak	<ul style="list-style-type: none">Light levels too highHigh day/night temperature above 70°F (21°C)
Chlorotic plants	<ul style="list-style-type: none">Media too wet – poor drainageHigh pH causing iron and nitrogen deficiencyAmmonium toxicityMagnesium deficiency

PanAmerican Seed™

PanAmSeed.com

PanAmerican Seed Co.
622 Town Road
West Chicago, Illinois USA 60185-2698
630 231-1400
Fax: 630 231-3609

PanAmerican Seed Europe BV
Lavendelweg 10
NL-1435 EW Rijsenhout, Holland
+31 (0)297-383038
Fax: +31 (0)297-383036

™ denotes a trademark of and ® denotes a registered trademark of Ball Horticultural Company in the U.S. It may also be registered in other countries.

©2005 Ball Horticultural Company Printed in USA PAS05161
PRIMLET is bred by and is a trademark of Kirin Brewery Co., Ltd. in the US.
It may also be registered in other countries.