

Grower Facts	Fantastic Flower class	Series/Variety	Recommended plug size**	Cover seed	Germination temperature	Days to germinate	406	288	128	Plug PGR recommendations	Comments
Pg 46	AGERATUM <i>A. houstonianum</i>	High Tide F1 Series	406-cell or larger	No	72-75°F (22-24°C)	3	4	4-5	5-6	No need	
Pg 48	ANGELONIA <i>A. angustifolia</i>	Serena™ F1 Series	406-cell or larger	No	72-76°F (22-25°C)	4-5	5-6	5-6	6-7	Generally no need. If necessary, B-Nine 5,000 ppm at Stage 3. Do not use growth regulator before radicle emergence as this can delay or stop germination.	Requires light to germinate. Grow plants on dry side, but do not allow plants to wilt.
Pg 49	BACOPA <i>Sutera cordata</i>	Blutopia™ Blue, Snowtopia™ White Improved	288 or 128-cell or larger	No	68-74°F (20-23°C)	4	—	3-4	4	B-Nine 2,500 ppm after transplant.	
Pg 57	DIASCIA <i>D. barberae</i>	Diamonte F1 Series	406-cell or larger	cover lightly	65-70°F (18-21°C)	4-6	4	4-5	—	B-Nine 2,500 ppm at Stage 3	Maintain media pH at 5.8-6.2.
Online*	HELENIUM <i>H. amarum</i>	Dakota Gold	288-cell or larger	Yes	65-75°F (18-22°C)	3-5	3-4	3-4	4-5	No need	Seedlings will grow slowly when daylength is shorter than 10 hours.
Pg 66	IMPATIENS, NEW GUINEA <i>I. hawkerii</i>	Divine F1 Series	288 to 128-cell	Yes	72-78°F (22-26°C)	6-8	5-6	5-6	6-7	No need	Supplemental lighting is not required, but will decrease total crop time.
Pg 68	LINARIA <i>L. hybrida</i>	Enchantment F1	406-cell or larger	Yes	65-68°F (18-20°C)	2-3	4-5	5-6	—	Bonzi 2 ppm at radicle emergence, then Bonzi 5 ppm spray 1 week later.	Germination is fast — 2-3 days. Make sure to move the seedling trays out of germination chamber on time to avoid stretch. Stage 1 PGR is key for avoiding leggy seedlings. Do not allow seedlings to wilt.
Pg 73	NEMESIA <i>N. foetans</i>	Poetry F1 Series	406-cell or larger	Yes	65-70°F (19-21°C)	4-5	4	4-5	—	B-Nine 2,500 ppm at Stage 3, if necessary. Do not use growth regulator before radicle emergence as this can delay or stop germination.	Seedlings may be more stretchy if germinating under dark conditions. Maintain media pH at 5.8-6.2.
Pg 78	PETUNIA, SMALL-FLOWERED SPREADING F1 <i>P. x hybrida</i>	Baby Duck Yellow	406-cell or larger	No	72-76°F (22-25°C)	4-7	4	5	6-7	B-Nine 5,000 ppm at Stage 3.	Baby Duck Yellow will flower successfully at 10 hours, however the crop time at 10 hours will be about 10-14 days longer than at 12-hour daylength.
Online*	SCUTELLARIA <i>S. javanica</i>	Veranda	406 or 288-cell	No	70-75°F (21-24°C)	6-10	6	7	—	No need	A heatloving crop — the warmer the temperature is, the faster the crop time. Crop time will be significantly longer at temperatures below 66°F (19°C).
Pg 94	TECOMA <i>T. stans</i>	Mayan Gold	288-cell or larger	Yes	68-74°F (20-23°C)	3-5	—	5-6	5-6	B-Nine 2,500 ppm and Cycocel 1,000 ppm mix every other week 2-3 weeks after transplanting.	
Pg 97	ZINNIA <i>Z. maritima</i>	Solcito	406 to 288-cell	No	70-75°F (21-24°C)	3-4	4-5	4-5	—		