GrowerFacts

Hypoestes Splash Select™ Series

(Hypoestes phyllostachya)

Potted Plants Culture (revised 02/25/20)

Plug crop time: 4 to 5 weeks Transplant to finish: 6 to 7 weeks

- · The grower's choice for mass market pot production, with excellent shipping tolerance and shelf life.
- Crop time can be reduced by planting multiple-sown young plants in each pot.
- Heat tolerant and shade tolerant.
- Can be used as an indoor plant or offered with other bedding annuals.
- The garden height for this plant is 10 to 18 in./25 to 46 cm; the height listed above is potted height.

General Information

Exposure	Bloom Season	Height	Spread	Spacing
Partial Sun	-	4-6 in. (10-15 cm)	12-14 in. (30-36 cm)	12-16 in. (30-41 cm)

Germination

Seed Form	Recommended Plug Size	Seeds/Cell	Plug Crop Weeks	Days from 50% to maximum germination	Initial Media pH/EC (1:2)	Cover Seed
RAW	288	1	4-5	2-3	5.5-6.0 pH 0.5-0.75 mmhos/cm	Yes

Plug Production

	Stage 1	Stage 2	Stage 3	Stage 4
Moisture	Level 4	Level 3-4	Level 3	Level 3
Temperature	70-75°F (21-24°C)	66-68°F (19-20°C)	66-68°F (19-20°C)	66-68°F (19-20°C)
Light	Optional			
Fertiliser		Less than 100 ppm N (Less than 0.7 EC)	100 to 175 ppm N (0.7 to 1.2 EC)	100 to 175 ppm N (0.7 to 1.2 EC)

Propagation Key Tips

Too much light can cause leaves to curl.

Growing on to Finish

Growing on Temperature	Target Media pH/EC (1:2)	Fertiliser	Daylength
(day) 65-75°F (18-24°C) (night) 62-65°F (17-18°C)	5.5-6.0 pH 1.0-1.5 mmhos/cm		-

Crop Scheduling

Container Size	Plugs/Pot	Crop Time	Season	PGR
Cell Pack	1 (ppp)	5-6 (weeks)	Spring	daminozide/chlormequat chloride tank mix 1,000/500 ppm Spray
4"/4.5"/Quart/10 cm	3-4 (ppp)	5-6 (weeks)	Spring	daminozide/chlormequat chloride tank mix 1,000/500 ppm Spray

Finishing Key Tips

Too much light can cause leaves to curl. Grow under low light conditions (400 to 500 f.c./4,000 to 5,000 Lux).

NOTE: Growers should use the information presented here as guidelines only. PanAmerican Seed recommends that growers conduct a trial of products under their own conditions. Crop times will vary depending on the climate, location, time of year, and greenhouse environmental conditions. It is the responsibility of the grower to confirm the treatment is available in their region as well as read and follow all the current label directions relating to the products. Nothing herein shall be deemed a warranty or guaranty by PanAmerican Seed of any products listed herein. PanAmerican Seed's terms and conditions of sale shall apply to all products listed herein.

Variety Pictures









PanAmerican Seed ®

PanAmerican Seed Co.
622 Town Road, West Chicago, Illinois, USA, 60185

+1 800-231-7065 PanAmSeed.com