

## Coleus, Premium Sun Chocolate Covered Cherry

Annuals Culture (revised 03/14/17)

*(Solenostemon scutellarioides)*

Show-stopping colours...bold patterns...rich textures: Pro landscapers and gardeners love these centre stage varieties for their great looks and reliable performance in sunny gardens and patio planters. Our Premium Sun Coleus can tolerate full sun in regions of high relative humidity such as Hardiness Zones 9 and 10. Foliage will scorch in full sun with low relative humidity. All are suitable for shade.

**Plug crop time:** 5 to 6 weeks

**Transplant to finish:** 6 to 8 weeks

- Boldly patterned, multi-coloured foliage makes a great companion to our other “chocolate” coleus: Chocolate Mint and Dark Chocolate.
- Non-fading colour pattern: rose centre on each leaf is surrounded by deep mahogany and edged with green.
- Late flowering and dwarf size are adaptable to pack and pot programs.
- Performs well in planters and small gardens in sun and shade.

### General Information

Exposure	Bloom Season	Height	Spread	Spacing
Sun	-	12-14 in. (30-36 cm)	10-12 in. (25-30 cm)	8 in. (20 cm)

### Germination

Seed Form	Recommended Plug Size	Seeds/Cell	Plug Crop Weeks	Days to Germinate	Initial Media pH/EC	Cover Seed
PEL	288	1	5-6	4-5	5.5-5.8 pH 0.75 mmhos/cm	Light cover

### Plug Production

	Stage 1	Stage 2	Stage 3	Stage 4
<b>Moisture</b>	Level 4	Level 3-4	Level 2-4	Level 2-4
<b>Temperature</b>	72-75°F (22-24°C)	72-75°F (22-24°C)	68-70°F (20-21°C)	60-62°F (16-17°C)
<b>Light</b>	Optional	2,500 f.c. (26,900 Lux)	2,500 f.c. (26,900 Lux)	2,500-5,000 f.c. (26,900-53,800 Lux)
<b>Fertilizer</b>	Less than 100 ppm N - Less than 0.7 EC		100 to 175 ppm N - 0.7 to 1.2 EC	100 to 175 ppm N - 0.7 to 1.2 EC

### Fertilizer Notation

To prevent diseases, irrigate early in the day so foliage is dry by nightfall.

### Propagation Key Tips

Coleus is very sensitive to high salts, particularly high ammonium. During germination, keep ammonium levels lower than 10 ppm. PGRs are generally not needed.

### Growing on to Finish

Growing on Temperature	Target Media pH/EC	Fertilizer	Daylength
<b>(day)</b> 68-75°F (20-24°C) <b>(night)</b> 65-68°F (18-20°C)	5.8-6.2 pH 1.5-2.0 mmhos/cm	100 to 175 ppm N - 0.7 to 1.2 EC	Facultative Short Day

## Crop Scheduling

Container Size	Plants/Pot	Crop Time	Season	PGR
306 Pack	1 (ppp)	6-8 (weeks)	Spring	-
4"/4.5"/Quart	1 (ppp)	6-8 (weeks)	Spring	-
5"/6"/1 Gallon	1 (ppp)	6-8 (weeks)	Spring	-
5"/6"/1 Gallon	3 (ppp)	6-8 (weeks)	Spring	-

### Common Problems

Insects: Aphids, Mealy Bugs, Whiteflies Diseases: Alternaria, Botrytis, Verticillium, Downy Mildew Other: Excessive internode elongation under low light or crowding.

### Finishing Key Tips

Excessive internode elongation under low light or crowding. Florel (ethephon) can be applied for promoting increased branching and height control if necessary. A rate of 300 ppm (7.69 ml/l 3.9% formulation or 0.63 ml/l of 48% formulation) at 2 to 3 weeks after transplanting is effective. Florel also delays flowering. Optional PGR: B-Nine/Alar (daminozide) 2,500 to 5,000 ppm (3.0 to 6.0 g/l 85% formulation or 4.0 to 8.0 g/l of 64% formulation) can be applied for height control at 2 to 3 weeks after transplanting. Repeat if necessary.

**NOTE:** Growers should use the information presented here as a starting point. Crop times will vary depending on the climate, location, time of year, and greenhouse environmental conditions. Chemical and PGR recommendations are only guidelines. It is the responsibility of the applicator to read and follow all the current label directions for the specific chemical being used in accordance with all regulations.

## Variety Pictures



Chocolate  
Covered Cherry



Chocolate  
Covered Cherry

PanAmericanSeed™

PanAmerican Seed Co.  
622 Town Road, West Chicago, Illinois, USA, 60185-2698  
630 231-1400 Fax: 630 231-3609 PanAmSeed.com

™ denotes a trademark of and ® denotes a registered trademark of Ball Horticultural Company in the US. It may also be registered in other countries.  
©2017 Ball Horticultural Company