

## Tomato Artemis F<sub>1</sub>

(*Solanum lycopersicum*)

Vegetables Culture (revised 01/19/21)

- One bite of these super-sweet, rich, red cherry tomatoes will have you craving more.
- The unique, crisp, firm texture is great in a salad or for sauce.
- High disease resistance to root-zone and foliar disease will ensure excellent plant health.
- Best grown in-ground, including high tunnels.

### General Information

Exposure	Bloom Season	Height	Spread	Spacing
Sun	Spring, Late Spring, Summer	60-84 in. (152-213 cm)	36-48 in. (91-122 cm)	18-36 in. (46-91 cm)

### Germination

Seed Form	Recommended Plug Size	Seeds/Cell	Plug Crop Weeks	Days from 50% to maximum germination	Initial Media pH/EC (1:2)	Cover Seed
RAW	288	1	3-4	2-3	5.5-5.8 pH 0.5 mmhos/cm	Light cover

### Plug Production

	Stage 1	Stage 2	Stage 3	Stage 4
<b>Moisture</b>	Level 4	Level 3-4	Level 2-3	Level 2-3
<b>Temperature</b>	70-75°F (21-24°C)	68-72°F (20-22°C)	68-72°F (20-22°C)	65-68°F (18-20°C)
<b>Light</b>	Optional	1,000-2,500 f.c. (10,800-26,900 Lux)	1,000-2,500 f.c. (10,800-26,900 Lux)	2,500-5,000 f.c. (26,900-53,800 Lux)
<b>Fertiliser</b>	Less than 100 ppm N (Less than 0.7 EC)	Less than 100 ppm N (Less than 0.7 EC)	Less than 100 ppm N (Less than 0.7 EC)	Less than 100 ppm N (Less than 0.7 EC)

### Fertiliser Notation

Tomatoes are very sensitive to high salts, particularly high ammonium, during germination. Keep ammonium level to less than 10 ppm.

### Propagation Key Tips

Height can be controlled by withholding fertilizer, avoiding overwatering, growing under high light conditions and using DIF when possible. If necessary, uniconazole (Sumagic) at a rate of 2.5 to 5 ppm can be applied at 2 weeks after sowing for height control. Repeat 2 weeks later if needed.

### Growing on to Finish

Growing on Temperature	Target Media pH/EC (1:2)	Fertiliser	Daylength
(day) 65-70°F (18-21°C) (night) 62-65°F (17-18°C)	5.5-6.2 pH 0.75-1.0 mmhos/cm	Less than 100 ppm N (Less than 0.7 EC)	-

### Days to Ripe Fruit

50-55

## Crop Scheduling

Container Size	Plugs/Pot	Crop Time	Season	PGR
Cell Pack	1 (ppp)	2-4 (weeks)	Spring	-
4"/4.5"/Quart/10 cm	1 (ppp)	2-4 (weeks)	Spring	-
Field grown	1 (ppp)	7-8 (weeks)	Spring	-

### Finishing Key Tips

Indeterminate variety. Performs best when grown in-ground and with support.

NOTE: Growers should use the information presented here as guidelines only. PanAmerican Seed recommends that growers conduct a trial of products under their own conditions. Crop times will vary depending on the climate, location, time of year, and greenhouse environmental conditions. It is the responsibility of the grower to confirm the treatment is available in their region as well as read and follow all the current label directions relating to the products. Nothing herein shall be deemed a warranty or guaranty by PanAmerican Seed of any products listed herein. PanAmerican Seed's terms and conditions of sale shall apply to all products listed herein.

## Variety Pictures



Artemis



PanAmerican Seed Co.  
622 Town Road, West Chicago, Illinois, USA, 60185  
+1 800-231-7065 [PanAmSeed.com](http://PanAmSeed.com)

™ denotes a trademark of and ® denotes a registered trademark of Ball Horticultural Company in the US. It may also be registered in other countries.  
©2024 Ball Horticultural Company