GrowerFacts

Herb (Basil) Monteverde Genovese

(Ocimum basilicum)

Vegetables Culture (revised 09/12/25)

Resists downy mildew for better harvests

Compared to other Genovese basils, NEW Monteverde Genovese lasts longer in the garden due to resistance to basil downy mildew. Added disease resistance means more reliable production of basil plants in all types of growing systems. Consumers will love the continual harvest.

- · Resistant to downy mildew and features a rich, Genovese basil flavour.
- Perfect for continual harvest throughout the season.
- Excellent for chopping fresh and adding to pasta dishes and salads, or for making pesto!

General Information

Exposure	Bloom Season	Height	Spread	Spacing
Sun	Late Spring, Summer, Late Summer	18-24 in. (46-61 cm)	10-12 in. (25-30 cm)	8-12 in. (20-30 cm)

Germination

Seed Form	Recommended Plug Size	Seeds/Cell	Plug Crop Weeks	Days to Germinate	Initial Media pH/EC	Cover Seed
RAW	288	6-10	3-5	2-4	5.5-5.8 pH 0.75 mmhos/cm	Yes

Plug Production

	Stage 1	Stage 2	Stage 3	Stage 4
Moisture	Level 4	Level 3-4	Level 2-3	Level 2-3
Temperature	68-74°F (20-23°C)	65-70°F (18-21°C)	65-70°F (18-21°C)	62-65°F (17-18°C)
Light	Optional	1,000-2,500 f.c. (10,800-26,900 Lux)	1,000-2,500 f.c. (10,800-26,900 Lux)	2,500-5,000 f.c. (26,900-53,800 Lux)
Fertiliser	Less than 100 ppm N (Less than 0.7 EC)	Less than 100 ppm N (Less than 0.7 EC)	Less than 100 ppm N (Less than 0.7 EC)	100 to 175 ppm N (0.7 to 1.2 EC)

Propagation Key Tips

Can be direct sown into final container.

Growing on to Finish

Growing on Temperature	Target Media pH/EC	Fertiliser	Daylength
(day) 65-70°F (18-21°C)	5.5-6.2 pH	100 to 175 ppm N	-
(night) 62-65°F (17-18°C)	1.0 mmhos/cm	(0.7 to 1.2 EC)	

Crop Scheduling

Container Size	Plugs/Pot	Crop Time	Season	PGR
Cell Pack	1 (ppp)	3-4 (weeks)	Spring	-
4"/4.5"/Quart/10 cm	1 (ppp)	3-4 (weeks)	Spring	-
5"/6"/1 Gallon/15 cm	1-3 (ppp)	4-5 (weeks)	Spring	-

Finishing Key Tips

Seed can be directly sown into finish containers.

NOTE: Growers should use the information presented here as guidelines only. PanAmerican Seed recommends that growers conduct a trial of products under their own conditions. Crop times will vary depending on the climate, location, time of year, and greenhouse environmental conditions. It is the responsibility of the grower to confirm the treatment is available in their region as well as read and follow all the current label directions relating to the products. Nothing herein shall be deemed a warranty or guaranty by PanAmerican Seed of any products listed herein. PanAmerican Seed's terms and conditions of sale shall apply to all products listed herein.

