

## Coreopsis Early Sunrise

(*Coreopsis grandiflora*)

**Plug crop time:** 5 to 7 weeks

**Transplant to finish:** 11 to 13 weeks

Perennials Culture (revised 10/14/25)

- Our most vigorous Coreopsis.
- Produces golden-yellow, semi-double flowers continuously all Summer.
- Can also be marketed as an annual in premium packs.
- For 2-gallon and larger containers in Summer, Early Sunrise is the optimal choice.

### General Information

Exposure	Bloom Season	Height	Spread	Spacing
Sun	Spring, Late Spring, Summer, Late Summer, Autumn	24 in. (61 cm)	22-24 in. (56-61 cm)	8-10 in. (20-25 cm)

### Germination

Seed Form	Recommended Plug Size	Seeds/Cell	Plug Crop Weeks	Days from 50% to maximum germination	Initial Media pH/EC (1:2)	Cover Seed
PRM	288 128*	1-2 2-4	5-7 7-8	3-5	5.8-6.2 pH 0.75 mmhos/cm	Light cover

\* Recommend plug size for forcing.

### Plug Production

	Stage 1	Stage 2	Stage 3	Stage 4
<b>Moisture</b>	Level 4	Level 3-4	Level 2-3	Level 1-2
<b>Temperature</b>	65-68°F (18-20°C)	65-68°F (18-20°C)	60-65°F (16-18°C)	60-65°F (16-18°C)
<b>Light</b>	Optional	6-8 mol·m <sup>-2</sup> ·d <sup>-1</sup>	8-10 mol·m <sup>-2</sup> ·d <sup>-1</sup>	8-10 mol·m <sup>-2</sup> ·d <sup>-1</sup>
<b>Fertiliser</b>		Less than 100 ppm N (Less than 0.7 EC)	100 to 175 ppm N (0.7 to 1.2 EC)	175 to 225 ppm N (1.2 to 1.5 EC)

### Vernalisation

No

### Propagation Key Tips

Critical Daylength: 14 hours.

For forcing info: See Perennials Forcing Guide.

1 to 2 seeds for 288/4 to 6 seeds for 84. Plug PGRs: For Spring production, no PGRs needed. For Summer production under long day conditions, can use daminozide (1,500 to 2,500 ppm spray). Short day bulk not needed for Autumn sales.

### Growing on to Finish

Growing on Temperature	Target Media pH/EC (1:2)	Fertiliser	Daylength
(day) 60-70°F (16-21°C) (night) 55-60°F (13-16°C)	5.8-6.2 pH 1.2-1.5 mmhos/cm	175 to 225 ppm N (1.2 to 1.5 EC)	Obligate Long Day

### Daylength Notation

Long day required – minimum 14 hours.

## Crop Scheduling

---

Crop Type	Container Size	Plugs/Pot	Crop Time	Season	PGR
Annual	4"/4.5"/Quart/10 cm	1 (ppp)	11-13 (weeks)	Summer	daminozide 5,000 ppm Spray
Annual	5"/6"/1 Gallon/15 cm	1-3 (ppp)	12-13 (weeks)	Summer	daminozide 5,000 ppm Spray
Forcing	5"/6"/1 Gallon/15 cm	1-3 (ppp)	12-13 (weeks)	Late Spring, ADT 68°F (20°C)	daminozide 2,500-5,000 ppm Spray
Forcing	5"/6"/1 Gallon/15 cm	1-3 (ppp)	11-12 (weeks)	Autumn, ADT 72°F (22°C)	daminozide 2,500-5,000 ppm Spray

## Common Problems

Insect: Aphid, Thrips, Whiteflies Disease: Powdery Mildew

## Bulking Strategy

SD bulk not needed for Early Sunrise.

## Finishing Key Tips

Apply PGRs when buds are visible.

NOTE: Growers should use the information presented here as guidelines only. PanAmerican Seed recommends that growers conduct a trial of products under their own conditions. Crop times will vary depending on the climate, location, time of year, and greenhouse environmental conditions. It is the responsibility of the grower to confirm the treatment is available in their region as well as read and follow all the current label directions relating to the products. Nothing herein shall be deemed a warranty or guaranty by PanAmerican Seed of any products listed herein. PanAmerican Seed's terms and conditions of sale shall apply to all products listed herein.

## Variety Pictures

---



Early Sunrise



Early Sunrise



PanAmerican Seed Co.  
622 Town Road, West Chicago, Illinois, USA, 60185  
+1 800-231-7065 PanAmSeed.com

™ denotes a trademark of and ® denotes a registered trademark of Ball Horticultural Company in the US. It may also be registered in other countries.  
©2026 Ball Horticultural Company