

Cyclamen Dreamscape™ F₁ Series

(*Cyclamen persicum*)

Annuals Culture (revised 02/24/20)

Plug crop time: 8 to 9 weeks

Transplant to finish: 19 to 20 weeks

- Specially bred and selected to deliver the best outdoor performance of any cyclamen.
- Pro landscapers and home gardeners alike can now confidently plant cyclamen in-ground and count on better, longer performance.
- Sow all colours at the same time to ship and sell together.

General Information

Exposure	Bloom Season	Height	Spread	Spacing
Shade, Partial Sun	Early Spring, Late Summer, Autumn, Winter	6-9 in. (15-23 cm)	9-13 in. (23-33 cm)	9-15 in. (23-38 cm)

Germination

Seed Form	Recommended Plug Size	Seeds/Cell	Plug Crop Weeks	Days to Germinate	Initial Media pH/EC	Cover Seed
RAW	288 128	1 1	11-12 14	19-21	5.6-5.8 pH 0.8 mmhos/cm	Yes

Plug Production

	Stage 1	Stage 2	Stage 3	Stage 4
Moisture	Level 4-5	Level 4-5	Level 3-4	Level 2-3
Temperature	64°F (18°C)	62-64°F (17-18°C)	61-64°F (16-18°C)	61-64°F (16-18°C)
Light	Dark	10-15 mol·m ⁻² ·d ⁻¹		
Fertilizer	Less than 100 ppm N (Less than 0.7 EC)	100 to 175 ppm N (0.7 to 1.2 EC)	100 to 175 ppm N (0.7 to 1.2 EC)	100 to 175 ppm N (0.7 to 1.2 EC)

Propagation Key Tips

Option is to transplant the plug into 60-72 cell tray and extend plug stage around 6-7 weeks before potting. This will reduce crop time after potting and plugs can be grown in a more controlled condition for a longer period.

Growing on to Finish

Growing on Temperature	Target Media pH/EC	Fertilizer	Daylength
(day) 62-64°F (17-18°C) (night) 54-59°F (12-15°C)	5.6-5.8 pH 0.8-1.2 mmhos/cm	225 to 300 ppm N (1.5 to 2.0 EC)	Day Neutral

Crop Scheduling

Container Size	Plugs/Pot	Crop Time	Season	PGR
5"/6"/1 Gallon	1 (ppp)	18-20 (weeks)	Autumn	-

Common Problems

Common Diseases: Botrytis, Chalara, Colletotrichum, Erwinia (bacterial soft rot), Fusarium oxysporum, Phytophthora, Pythium, Rhizoctonia, Cylindrocarpum
Common Pests: Aphids, caterpillars (Duponchelia), Thrips, Cyclamen mites, Fungus gnats and Shore Flies

Finishing Key Tips

Use of 80-72 cell plugs will reduce crop time after potting for around 6-7 weeks.

NOTE: Growers should use the information presented here as guidelines only. PanAmerican Seed recommends that growers conduct a trial of products under their own conditions. Crop times will vary depending on the climate, location, time of year, and greenhouse environmental conditions. It is the responsibility of the grower to confirm the treatment is available in their region as well as read and follow all the current label directions relating to the products. Nothing herein shall be deemed a warranty or guaranty by PanAmerican Seed of any products listed herein. PanAmerican Seed's terms and conditions of sale shall apply to all products listed herein.

Variety Pictures



Bright Red



Deep Burgundy



Deep Rose



Purple



Salmon Eye



White



Dark Flamed
Mixture



Mixture

PanAmericanSeed™

PanAmerican Seed Co.
622 Town Road, West Chicago, Illinois, USA, 60185-2698
630 231-1400 Fax: 630 231-3609 PanAmSeed.com

™ denotes a trademark of and © denotes a registered trademark of Ball Horticultural Company in the US. It may also be registered in other countries.
©2021 Ball Horticultural Company