

## Lavandula Ellagance Series

(*Lavandula angustifolia*)

Perennials Culture (revised 02/25/20)

**Plug crop time:** 5 to 6 weeks

**Transplant to finish:** 10 to 14 weeks (24 to 36 weeks overwintered)

- Ellagance stands out from other seed and vegetative lavenders for its excellent uniformity, colour range, earliness and spike quality.
- Outstanding branching fills pots quickly and finishes with large, dense flower spikes and very strong stems.
- Spreading and upright plants have rich grey-green foliage.
- Winter hardy, cold tolerant and durable performance; foliage holds up better compared to other lavenders.
- High uniformity means harvest times are short and comparable to any vegetative variety.
- Ships well; long-flowering in perennial beds and patio planters.
- Great as a garden plant and for drying and arranging.
- Large, elegant flowers bloom all Summer and attract butterflies.

### General Information

Exposure	Bloom Season	Height	Spread	Spacing
Sun	-	12-14 in. (30-36 cm)	10-12 in. (25-30 cm)	8-12 in. (20-30 cm)

### Germination

Seed Form	Recommended Plug Size	Seeds/Cell	Plug Crop Weeks	Days to Germinate	Initial Media pH/EC	Cover Seed
PRM	288 128*	4 6	6-8 8-9	4-5	5.8-6.5 pH 0.7-1.2 mmhos/cm	Yes

\* Recommend plug size for forcing.

### Plug Production

	Stage 1	Stage 2	Stage 3	Stage 4
<b>Moisture</b>	Level 4	Level 3	Level 2-3	Level 2
<b>Temperature</b>	65-68°F (18-20°C)	65-68°F (18-20°C)	60-65°F (16-18°C)	60-65°F (16-18°C)
<b>Light</b>	Light	2,000-2,500 f.c. (21,500-26,900 Lux)	2,500-3,500 f.c. (26,900-37,700 Lux)	4,000-5,000 f.c. (43,100-53,800 Lux)
<b>Fertilizer</b>	100 to 175 ppm N - 0.7 to 1.2 EC	100 to 175 ppm N - 0.7 to 1.2 EC	175 to 225 ppm N - 1.2 to 1.5 EC	175 to 225 ppm N - 1.2 to 1.5 EC
<b>PGR</b>			daminozide/1,000-2,000 ppm/Spray	daminozide/1,000-2,000 ppm/Spray

### Vernalization

No

### Propagation Key Tips

For forcing info: see Perennials Forcing Guide.

For scheduling info: see Lavender scheduling tool.

Spray damp-off fungicide. Provide good ventilation and active respiration in plug production. Grow on the dry side, with watering in early morning, to allow plugs to dry up during the day. If respiration and fertilization are too low, *Lavandula angustifolia* can show "black spots" on cotyledons due to root pressure damage. This is reversible with improved ventilation and higher fertilization.

## Growing on to Finish

---

Growing on Temperature	Target Media pH/EC	Fertilizer	Daylength
(day) 60-72°F (16-22°C) (night) 54-60°F (12-16°C)	5.8-6.5 pH 1.1-1.4 mmhos/cm	100 to 175 ppm N - 0.7 to 1.2 EC	Facultative Long Day

### Daylength Notation

Critical Daylength for Ellagance Purple and Pink is 10 hours; all other varieties are Obligate Long Day with critical daylength of about 13-14 hours.

See Lavender Scheduling Tool at panamseed.com for finishing schedules by region.

## Crop Scheduling

---

Crop Type	Container Size	Plugs/Pot	Crop Time	Season	PGR
Overwinter	4"/4.5"/Quart	1 (ppp)	28-36 (weeks)	Spring, ADT 65°F (18°C)	daminozide 2,000-3,000 ppm Spray
Annual	4"/4.5"/Quart	1 (ppp)	9-12 (weeks)	Late Spring, ADT 65°F (18°C)	daminozide 2,000-3,000 ppm Spray
Annual	5"/6"/1 Gallon	1-3 (ppp)	9-12 (weeks)	Late Spring, ADT 65°F (18°C)	daminozide 2,000-3,000 ppm Spray

### Fertilizer Notation

Apply constant fertilizer at rate 1 to 2 (100 to 125 ppm N/0.7 to 1.2 mmhos/cm EC or 150 ppm as needed). Maintain the pH at 5.8 to 6.5 and EC 1.2 to 1.5 mmhos/cm. For overwintering outside, add some extra nitrate-based fertilizer in early Spring (EC 1.5 to 1.6 mmhos/cm) when plants start to re-grow.

### Common Problems

Needs active climate, with good airflow, low RH and grow relatively dry. Must have excellent soil drainage, but also container field drainage. Will be prone to fungal disease (Leafspot, Botrytis) when high RV, little airflow, low light. Lavandula plants need to be dry when being shipped. No overhead watering in shipping, nor at retail environment. For Long day Summer production, try Lavance Deep Purple; this has better bulk and stronger color during long and high-light Summer days.


### Finishing Key Tips

Avoid planting plugs deep. Keep plug surface same as medium surface. Grow low RH and high light. Let top soil dry in between waterings, but do not allow medium to dry as high EC could cause root damage. Monitor for Botrytis, Root Rot, Leafspot, Aphids and Mites. For shipping, keep soil moist and plant dry. Scheduling is different between colours: Purple and Pink are the same and the fastest (as indicated lead time); Sky adds 1 week; Ice adds 3 weeks, and Snow adds 4 to 5 weeks. See Scheduling Tool for regional lead times. For overwintering production, cut back just below top foliage; do this mid-Autumn, approximately 4 weeks before dormant period, or in Spring just before regrowth.

NOTE: Growers should use the information presented here as guidelines only. PanAmerican Seed recommends that growers conduct a trial of products under their own conditions. Crop times will vary depending on the climate, location, time of year, and greenhouse environmental conditions. It is the responsibility of the grower to confirm the treatment is available in their region as well as read and follow all the current label directions relating to the products. Nothing herein shall be deemed a warranty or guaranty by PanAmerican Seed of any products listed herein. PanAmerican Seed's terms and conditions of sale shall apply to all products listed herein.

## Variety Pictures


---



Ice



Purple



Snow

PanAmericanSeed™

PanAmerican Seed Co.  
622 Town Road, West Chicago, Illinois, USA, 60185-2698  
**630 231-1400 Fax: 630 231-3609 PanAmSeed.com**

™ denotes a trademark of and © denotes a registered trademark of Ball Horticultural Company in the US. It may also be registered in other countries.  
©2020 Ball Horticultural Company