# Lavandula Ellagance Series

(Lavandula angustifolia)

Perennials Culture (revised 12/02/24)

#### Complete your Lavender program here!

Find four outstanding varieties to fill in your regionally focused program of *Lavandula angustifolia* from PanAmerican Seed. With Blue Spear, Avignon Early Blue, Ellagance Purple and Lavance Deep Purple, we have all the seed Lavender to keep you growing. Find detailed information in our regional Lavender Scheduling Tool at panamseed.com/perennials.

Plug crop time: 5 to 6 weeks

Transplant to finish: 10 to 14 weeks (24 to 36 weeks overwintered)

- · Ellagance stands out from other seed and vegetative lavenders for its excellent uniformity, earliness and spike quality.
- · Outstanding branching fills pots quickly and finishes with large, dense flower spikes and very strong stems.
- Spreading and upright plants have rich grey-green foliage.
- Winter hardy, cold tolerant and durable performance; foliage holds up better compared to other lavenders.
- · High uniformity means harvest times are short and comparable to any vegetative variety.
- Ships well; long-flowering in perennial beds and patio planters.
- · Great as a garden plant and for drying and arranging.
- · Large, elegant flowers bloom all Summer and attract butterflies.

### **General Information**

| Exposure | Bloom Season | Height                  | Spread                  | Spacing                |
|----------|--------------|-------------------------|-------------------------|------------------------|
| Sun      | Summer       | 12-14 in.<br>(30-36 cm) | 10-12 in.<br>(25-30 cm) | 8-12 in.<br>(20-30 cm) |

### **Germination**

| Seed Form | Recommended<br>Plug Size | Seeds/Cell | Plug Crop Weeks | Days from 50% to maximum germination | Initial Media<br>pH/EC (1:2)   | Cover Seed |
|-----------|--------------------------|------------|-----------------|--------------------------------------|--------------------------------|------------|
| PRM       | 288<br>128*              | 4<br>6     | 6-8<br>8-9      | 4-5                                  | 5.8-6.5 pH<br>0.7-1.2 mmhos/cm | Yes        |

<sup>\*</sup> Recommend plug size for forcing.

# **Plug Production**

|             | Stage 1              | Stage 2                                 | Stage 3                                 | Stage 4                                 |
|-------------|----------------------|---|---|---|
| Moisture    | Level 4              | Level 3                                 | Level 2-3                               | Level 2                                 |
| Temperature | 65-68°F<br>(18-20°C) | 65-68°F<br>(18-20°C)                    | 60-65°F<br>(16-18°C)                    | 60-65°F<br>(16-18°C)                    |
| Light       | Light                | 2,000-2,500 f.c.<br>(21,500-26,900 Lux) | 2,500-3,500 f.c.<br>(26,900-37,700 Lux) | 4,000-5,000 f.c.<br>(43,100-53,800 Lux) |
| Fertiliser  |                      | 100 to 175 ppm N<br>(0.7 to 1.2 EC)     | 175 to 225 ppm N<br>(1.2 to 1.5 EC)     | 175 to 225 ppm N<br>(1.2 to 1.5 EC)     |
| PGR         |                      |   | daminozide/1,000-2,000 ppm/Spray        | daminozide/1,000-2,000 ppm/Spray        |

#### Vernalisation

No

### **Propagation Key Tips**

For forcing info: see Perennials Forcing Guide. For scheduling info: see Lavender scheduling tool.

Spray damp-off fungicide. Provide good ventilation and active respiration in plug production. Grow on the dry side, with watering in early morning, to allow plugs to dry up during the day. If respiration and fertilisation are too low, Lavandula angustifolia can show "black spots" on cotyledons due to root pressure damage. This is reversible with improved ventilation and higher fertilisation.

### **Growing on to Finish**

| Growing on<br>Temperature | Target Media<br>pH/EC (1:2) | Fertiliser       | Daylength            |
|---------------------------|-----------------------------|------------------|----------------------|
| (day) 60-72°F (16-22°C)   | 5.8-6.5 pH                  | 100 to 175 ppm N | Facultative Long Day |
| (night) 54-60°F (12-16°C) | 1.1-1.4 mmhos/cm            | (0.7 to 1.2 EC)  |                      |

### **Daylength Notation**

Critical Daylength for Ellagance Purple is 10 hours; Ice and Snow are Obligate Long Day with critical daylength of about 13-14 hours. See Lavender Scheduling Tool at panamseed.com for finishing schedules by region.

### **Crop Scheduling**

| Crop Type  | Container Size       | Plugs/Pot | Crop Time     | Season                       | PGR                              |
|------------|----------------------|-----------|---------------|------------------------------|----------------------------------|
| Overwinter | 4"/4.5"/Quart/10 cm  | 1 (ppp)   | 28-36 (weeks) | Spring, ADT 65°F (18°C)      | daminozide 2,000-3,000 ppm Spray |
| Annual     | 4"/4.5"/Quart/10 cm  | 1 (ppp)   | 9-12 (weeks)  | Late Spring, ADT 65°F (18°C) | daminozide 2,000-3,000 ppm Spray |
| Annual     | 5"/6"/1 Gallon/15 cm | 1-3 (ppp) | 9-12 (weeks)  | Late Spring, ADT 65°F (18°C) | daminozide 2,000-3,000 ppm Spray |

#### **Fertiliser Notation**

Apply constant fertilizer at rate 1 to 2 (100 to 125 ppm N/0.7 to 1.2 mmhos/cm EC or 150 ppm as needed). Maintain the pH at 5.8 to 6.5 and EC 1.2 to 1.5 mmhos/cm. For overwintering outside, add some extra nitrate-based fertilizer in early Spring (EC 1.5 to 1.6 mmhos/cm) when plants start to re-grow.

#### **Common Problems**

Needs active climate, with good airflow, low RH and grow relatively dry. Must have excellent soil drainage, but also container field drainage. Will be prone to fungal disease (Leafspot, Botrytis) when high RV, little airflow, low light. Lavandula plants need to be dry when being shipped. No overhead watering in shipping, nor at retail environment. For Long day Summer production, try Lavance Deep Purple; this has better bulk and stronger color during long and high-light Summer days.

#### **Finishing Key Tips**

Avoid planting plugs deep. Keep plug surface same as medium surface. Grow low RH and high light. Let top soil dry in between waterings, but do not allow medium to dry as high EC could cause root damage. Monitor for Botrytis, Root Rot, Leafspot, Aphids and Mites. For shipping, keep soil moist and plant dry. Scheduling is different between colours: Purple is the fastest (as indicated lead time) and Snow adds 4 to 5 weeks. See Scheduling Tool for regional lead times. For overwintering production, cut back just below top foliage; do this mid-Autumn, approximately 4 weeks before dormant period, or in Spring just before regrowth.

NOTE: Growers should use the information presented here as guidelines only. PanAmerican Seed recommends that growers conduct a trial of products under their own conditions. Crop times will vary depending on the climate, location, time of year, and greenhouse environmental conditions. It is the responsibility of the grower to confirm the treatment is available in their region as well as read and follow all the current label directions relating to the products. Nothing herein shall be deemed a warranty or guaranty by PanAmerican Seed of any products listed herein. PanAmerican Seed's terms and conditions of sale shall apply to all products listed herein.

## **Variety Pictures**



