

Marigold (French) Janie Series

(*Tagetes patula*)

Annuals Culture (revised 10/23/18)

Plug crop time: 3 weeks

Transplant to finish: 3 to 4 weeks

- Good for warmer southern markets or for late-season sales in the North – very heat-tolerant with good garden vigour in warmer temperatures.
- Early and uniform in the pack with excellent shelf life.
- Compact plants hold crested, 2-in./5-cm flowers nicely above the foliage.

General Information

Exposure	Bloom Season	Height	Spread	Spacing
Sun	Spring, Late Spring, Summer, Late Summer	8-10 in. (20-25 cm)	6-8 in. (15-20 cm)	6-10 in. (15-25 cm)

Germination

Seed Form	Recommended Plug Size	Seeds/Cell	Plug Crop Weeks	Days to Germinate	Initial Media pH/EC	Cover Seed
COT, DTL	288	1	3-4	2-3	6.2-6.5 pH 0.75 mmhos/cm	Yes

Plug Production

	Stage 1	Stage 2	Stage 3	Stage 4
Moisture	Level 4	Level 3-4	Level 3	Level 3
Temperature	70-72°F (21-22°C)	68-72°F (20-22°C)	68-75°F (20-24°C)	68-75°F (20-24°C)
Light	Optional			
Fertilizer	Less than 100 ppm N - Less than 0.7 EC	Less than 100 ppm N - Less than 0.7 EC	100 to 175 ppm N - 0.7 to 1.2 EC	100 to 175 ppm N - 0.7 to 1.2 EC

Propagation Key Tips

Maintain soil pH of 6.2 to 6.5 to avoid iron toxicity.

Growing on to Finish

Growing on Temperature	Target Media pH/EC	Fertilizer	Daylength
(day) 65-68°F (18-20°C) (night) 60-62°F (16-17°C)	6.2-6.5 pH 1.0-1.5 mmhos/cm	100 to 175 ppm N - 0.7 to 1.2 EC	Facultative Short Day

Crop Scheduling

Container Size	Plugs/Pot	Crop Time	Season	PGR
Cell Pack	1 (ppp)	3-4 (weeks)	Spring	-

Fertilizer Notation

Starting 1 week after transplant, apply fertilizer at rate of 100 to 175 ppm N and 0.7 to 1.2 mS/cm using predominantly nitrate-based fertilizer with low phosphorus. Maintain the media EC at 1.0 to 1.5 mS/cm and pH at 6.2 to 6.5. For constant fertilizer program, fertilizer can be applied at a rate of 75 to 100 ppm N 0.5 to 0.7 mS/cm while maintaining the above recommended EC and pH ranges.

Common Problems

Insects: Aphids, Mites, Whiteflies Diseases: Damping off in the seedling stage

Finishing Key Tips


Maintain soil pH of 6.2 to 6.5 to avoid iron toxicity.

NOTE: Growers should use the information presented here as guidelines only. PanAmerican Seed recommends that growers conduct a trial of products under their own conditions. Crop times will vary depending on the climate, location, time of year, and greenhouse environmental conditions. It is the responsibility of the grower to read and follow all the current label directions relating to the products. Nothing herein shall be deemed a warranty or guaranty by PanAmerican Seed of any products listed herein. PanAmerican Seed's terms and conditions of sale shall apply to all products listed herein.

Variety Pictures



Bright Yellow



Deep Orange



Spry

PanAmericanSeed™

PanAmerican Seed Co.
622 Town Road, West Chicago, Illinois, USA, 60185-2698
630 231-1400 Fax: 630 231-3609 PanAmSeed.com

™ denotes a trademark of and ® denotes a registered trademark of Ball Horticultural Company in the US. It may also be registered in other countries.
©2018 Ball Horticultural Company