

Petunia Madness® F₁ Series

(*Petunia x hybrida*)

Annuals Culture (revised 10/30/24)

Plug crop time: 4 to 5 weeks

Transplant to finish: 5 to 6 weeks

Available in North America only.

- Home gardeners and pro landscapers across North America know they can count on fantastic flower power...a bounty of vibrant colours... and never-ending shows that last until first frost.
- Exceptional seed quality, plug performance and uniformity across all colours from sow to finish.
- Well-branched, floriferous plants are covered with an abundance of 3-in./8-cm blooms.

General Information

Exposure	Bloom Season	Height	Spread	Spacing
Sun	Spring, Late Spring, Summer, Late Summer	10-15 in. (25-38 cm)	10-12 in. (25-30 cm)	8-10 in. (20-25 cm)

Germination

Seed Form	Recommended Plug Size	Seeds/Cell	Plug Crop Weeks	Days from 50% to maximum germination	Initial Media pH/EC (1:2)	Cover Seed
PEL, RAW	288	1	4-5	3-5	5.5-5.8 pH 0.75-1.0 mmhos/cm	No

Plug Production

	Stage 1	Stage 2	Stage 3	Stage 4
Moisture	Level 4-5	Level 3-4	Level 2-4	Level 2-4
Temperature	72-77°F (22-25°C)	72-77°F (22-25°C)	65-70°F (18-21°C)	62-65°F (17-18°C)
Light	Light	1,000-2,500 f.c. (10,800-26,900 Lux)	2,500-4,000 f.c. (26,900-43,100 Lux)	2,500-5,000 f.c. (26,900-53,800 Lux)
Fertiliser		Less than 100 ppm N (Less than 0.7 EC)	100 to 175 ppm N (0.7 to 1.2 EC)	100 to 175 ppm N (0.7 to 1.2 EC)
PGR			daminozide/2,500-5,000 ppm/Spray	daminozide/2,500-5,000 ppm/Spray

Growing on to Finish

Growing on Temperature	Target Media pH/EC (1:2)	Fertiliser	Daylength
(day) 62-70°F (17-21°C) (night) 55-65°F (13-18°C)	5.5-6.2 pH 1.0-1.5 mmhos/cm	100 to 175 ppm N (0.7 to 1.2 EC)	Facultative Long Day

Crop Scheduling

Container Size	Plugs/Pot	Crop Time	Season	PGR
Cell Pack	1 (ppp)	5-6 (weeks)	Spring	daminozide 2,500-5,000 ppm Spray
10" Pot or HB/3 Gallon/25 cm	3 (ppp)	5-7 (weeks)	Spring	daminozide 2,500-5,000 ppm Spray

Common Problems


Insects: Thrips, Aphids, Fungus Gnats, Shore Flies Diseases: Botrytis, Rhizoctonia, Pythium

NOTE: Growers should use the information presented here as guidelines only. PanAmerican Seed recommends that growers conduct a trial of products under their own conditions. Crop times will vary depending on the climate, location, time of year, and greenhouse environmental conditions. It is the responsibility of the grower to confirm the treatment is available in their region as well as read and follow all the current label directions relating to the products. Nothing herein shall be deemed a warranty or guaranty by PanAmerican Seed of any products listed herein. PanAmerican Seed's terms and conditions of sale shall apply to all products listed herein.


Variety Pictures




Blue Star




Blue Vein




Burgundy



Burgundy Star




Deep Rose




Lilac




Mid-Blue




Midnight




Orchid




Pink




Plum Crazy




Red




Red Morn




Red Picotee




Rose




Rose Morn




Rose Star



Royal




Sheer



Simply




Sky Blue



Spring




Sugar



Summer



White



Yellow



Clear Mixture



Just Mixture



Merlot Mixture



Old Glory
Mixture



Total Mixture



Waterfall
Mixture



PanAmerican Seed Co.
622 Town Road, West Chicago, Illinois, USA, 60185
+1 800-231-7065 PanAmSeed.com

™ denotes a trademark of and © denotes a registered trademark of Ball Horticultural Company in the US. It may also be registered in other countries.
©2026 Ball Horticultural Company