

Primula Heritage Crème F₁

(*Primula acaulis*)

Annuals Culture (revised 03/16/17)

Plug crop time: 5 to 6 weeks

Transplant to finish: 15 to 17 weeks

- Unique breeding delivers a natural “English Primrose” look in a professional-quality product.
- Long-flowering, mid-size blooms and deep green foliage are produced on compact, mounded plants.
- Programmable crop can be grown pot-tight in 3.5-in./9-cm programs.

General Information

Exposure	Bloom Season	Height	Spread	Spacing
Partial Sun, Sun	Early Spring, Spring, Autumn	5-6 in. (13-15 cm)	5-6 in. (13-15 cm)	5-6 in. (13-15 cm)

Germination

Seed Form	Recommended Plug Size	Seeds/Cell	Plug Crop Weeks	Days to Germinate	Initial Media pH/EC	Cover Seed
RAW	288	1	5-6	7-10	5.5-5.8 pH 0.75 mmhos/cm	Light cover

Plug Production

	Stage 1	Stage 2	Stage 3	Stage 4
Moisture	Level 4-5	Level 3-4	Level 3-4	Level 3
Temperature	64-68°F (18-20°C)	60-62°F (16-17°C)	66-72°F (19-22°C)	64-68°F (18-20°C)
Light	Optional	1,000-2,500 f.c. (10,800-26,900 Lux)	1,000-2,500 f.c. (10,800-26,900 Lux)	2,500-5,000 f.c. (26,900-53,800 Lux)
Fertilizer	Less than 100 ppm N - Less than 0.7 EC	Less than 100 ppm N - Less than 0.7 EC	100 to 175 ppm N - 0.7 to 1.2 EC	100 to 175 ppm N - 0.7 to 1.2 EC

Propagation Key Tips

Avoid too high light levels (> 3,000 f.c.) to prevent leaf damage.

Growing on to Finish

Growing on Temperature	Target Media pH/EC	Fertilizer	Daylength
(day) 55-60°F (13-16°C) (night) 50-55°F (10-13°C)	5.5-6.2 pH 1.0-1.2 mmhos/cm	100 to 175 ppm N - 0.7 to 1.2 EC	Facultative Long Day

Crop Scheduling

Container Size	Plugs/Pot	Crop Time	Season	PGR
5"/6"/1 Gallon	1 (ppp)	15-17 (weeks)	Autumn	daminozide 2,500-5,000 ppm Spray

Fertilizer Notation

Start the plants with 20-10-20 at 200 ppm. When ready for cold treatment, change fertilizer to 15-0-15 at 50 ppm.

Common Problems

Insect: Aphids, Whiteflies, Fungus Gnats Disease: Botrytis on flowers, Pythium, Rhizoctonia

Finishing Key Tips

Drop temperature after week 5 (when plant has 10 established leaves) to 45 to 48°F (7 to 9°C) day and 35 to 45°F (2° to 7°C) night for bud initiation. After week 11, go back up to growing temperatures for flower development and forcing. Plants can be held at 40 to 45°F (5 to 7°C) for later forcing. Growing-on time in weeks depends on how large a plant is required. A large plant requires a longer time at 60 to 65°F (16 to 18°C) nights. From bud visibility to first opening of flower is approximately 4 to 5 weeks, depending on temperature.

NOTE: Growers should use the information presented here as guidelines only. PanAmerican Seed recommends that growers conduct a trial of products under their own conditions. Crop times will vary depending on the climate, location, time of year, and greenhouse environmental conditions. It is the responsibility of the grower to read and follow all the current label directions relating to the products. Nothing herein shall be deemed a warranty or guaranty by PanAmerican Seed of any products listed herein. PanAmerican Seed's terms and conditions of sale shall apply to all products listed herein.

Variety Pictures



Heritage Crème



Heritage Crème

PanAmericanSeed™

PanAmerican Seed Co.
622 Town Road, West Chicago, Illinois, USA, 60185-2698
630 231-1400 Fax: 630 231-3609 PanAmSeed.com

™ denotes a trademark of and ® denotes a registered trademark of Ball Horticultural Company in the US. It may also be registered in other countries.
©2018 Ball Horticultural Company