

Nepeta Pink Cat

(*Nepeta nervosa*)

Perennials Culture (revised 01/10/18)

Plug crop time: 5 to 6 weeks

Transplant to finish: 8 to 10 weeks

- Uniform strains produce large, dense flowers spikes on fully, bushy plants that flower early and then freely all season.
- Well-suited to container production.
- Gives a colourful display in well-drained soil and takes the heat.

General Information

Exposure	Bloom Season	Height	Spread	Spacing
Sun	Summer	10-12 in. (25-30 cm)	10-12 in. (25-30 cm)	

Germination

Seed Form	Recommended Plug Size	Seeds/Cell	Plug Crop Weeks	Days to Germinate	Initial Media pH/EC	Cover Seed
RAW	288	4	5-6	4-5	5.8-6.5 pH 0.75 mmhos/cm	Yes

Plug Production

	Stage 1	Stage 2	Stage 3	Stage 4
Moisture	Level 4	Level 3	Level 2	Level 2
Temperature	65-68°F (18-20°C)	60-65°F (16-18°C)	60-65°F (16-18°C)	60-65°F (16-18°C)
Light	Optional	2,000-2,500 f.c. (21,500-26,900 Lux)	2,000-2,500 f.c. (21,500-26,900 Lux)	4,000-5,000 f.c. (43,100-53,800 Lux)
Fertilizer	Less than 100 ppm N - Less than 0.7 EC	100 to 175 ppm N - 0.7 to 1.2 EC	100 to 175 ppm N - 0.7 to 1.2 EC	100 to 175 ppm N - 0.7 to 1.2 EC
PGR			daminozide/1,000-2,000 ppm/Spray	daminozide/1,500-2,000 ppm/Spray

Vernalization

No

Propagation Key Tips

Spray damp-off fungicide.

Growing on to Finish

Growing on Temperature	Target Media pH/EC	Fertilizer	Daylength
(day) 60-65°F (16-18°C) (night) 54-60°F (12-16°C)	5.8-6.5 pH 1.2-1.4 mmhos/cm	175 to 225 ppm N - 1.2 to 1.5 EC	Facultative Long Day

Daylength Notation

Long day beneficial.

Crop Scheduling

Crop Type	Container Size	Plugs/Pot	Crop Time	Season	PGR
Annual	4"/4.5"/Quart	1 (ppp)	8-12 (weeks)	Late Spring	daminozide 2,500 ppm Spray
Annual	5"/6"/1 Gallon	2-3 (ppp)	8-12 (weeks)	Late Spring	daminozide 2,500 ppm Spray

Finishing Key Tips

Grow relatively dry. Prevent Mg and Fe deficiencies. Monitor for Botrytis, Downy Mildew and Aphids.

NOTE: Growers should use the information presented here as guidelines only. PanAmerican Seed recommends that growers conduct a trial of products under their own conditions. Crop times will vary depending on the climate, location, time of year, and greenhouse environmental conditions. It is the responsibility of the grower to confirm the treatment is available in their region as well as read and follow all the current label directions relating to the products. Nothing herein shall be deemed a warranty or guaranty by PanAmerican Seed of any products listed herein. PanAmerican Seed's terms and conditions of sale shall apply to all products listed herein.

Variety Pictures



Pink Cat

PanAmericanSeed™

PanAmerican Seed Co.
622 Town Road, West Chicago, Illinois, USA, 60185-2698
630 231-1400 Fax: 630 231-3609 PanAmSeed.com

™ denotes a trademark of and ® denotes a registered trademark of Ball Horticultural Company in the US. It may also be registered in other countries.
©2019 Ball Horticultural Company