

Begonia (Tuberous) AmeriHybrid® Picotee F₁ Series

Annuals Culture (revised 02/14/23)

(*Begonia x tuberosa*)

Big, bold begonias

AmeriHybrid tuberous begonias provide a wide selection of colours, including unique bicolours, along with some of the world's largest blooms! This exciting collection can be sold at a premium price point at retail, and it results in faster turnover in the market.

Plug crop time: 7 to 8 weeks

Transplant to finish: 11 to 15 weeks

- The AmeriHybrid Picotee varieties have large, 4 to 8-in./10 to 20-cm, bicolour blooms.
- The Picotee Lace types are an inverted variety of a normal picotee; these blooms have light edges on dark-coloured petals.

General Information

| Exposure | Bloom Season | Height | Spread | Spacing |
|--------------------|-----------------------------|-------------------------|-------------------------|-------------------------|
| Shade, Partial Sun | Late Spring, Summer, Autumn | 14-20 in. (36-51 cm) | 10-12 in. (25-30 cm) | 12-16 in. (30-41 cm) |

Germination

| Seed Form | Recommended Plug Size | Seeds/Cell | Plug Crop Weeks | Days from 50% to maximum germination | Initial Media pH/EC (1:2) | Cover Seed |
|-----------|-----------------------|------------|-----------------|--------------------------------------|----------------------------|------------|
| PEL | 288 | 1 | 7-8 | 7-14 | 5.5-6.0 pH 0.5 mmhos/cm | No |

Plug Production

| | Stage 1 | Stage 2 | Stage 3 | Stage 4 |
|--------------------|----------------------|---|---|---|
| Moisture | Level 5 | Level 4-5 | Level 3-4 | Level 3-4 |
| Temperature | 72-76°F (22-24°C) | 68-72°F (20-22°C) | 65-68°F (18-20°C) | 62-68°F (17-20°C) |
| Light | Light | 350-600 f.c. (3,800-6,500 Lux) | 350-600 f.c. (3,800-6,500 Lux) | 500-1,000 f.c. (5,400-10,800 Lux) |
| Fertiliser | | Less than 100 ppm N (Less than 0.7 EC) | Less than 100 ppm N (Less than 0.7 EC) | Less than 100 ppm N (Less than 0.7 EC) |

Fertiliser Notation

Avoid high salt levels in the substrate, as it can cause leaf burn.

Propagation Key Tips

Keep moisture high until the first true leaf develops. Once cotyledons are visible, maintain moist but not saturated media to promote root development. Avoid using ammonium nitrate during plug development, as it may inhibit root growth. A minimum 14-hour photoperiod lighting is required to avoid tuber formation and improve quality. Avoid strong sunlight (>2,000 foot-candles). Strong sunlight will cause high leaf temperature and result in burned leaf edges.

Growing on to Finish

| Growing on Temperature | Target Media pH/EC (1:2) | Fertiliser | Daylength |
|--|--------------------------------|-------------------------------------|-------------------|
| (day) 70-72°F (21-22°C) (night) 62-68°F (17-20°C) | 5.5-5.8 pH 1.0-1.5 mmhos/cm | 175 to 225 ppm N (1.2 to 1.5 EC) | Obligate Long Day |

Daylength Notation

Maintain minimum of 14-hour photoperiod of light for active growth. A photoperiod of less than 12 hours will result in small single flowers and an uneven crop, and plants will form tubers.

Crop Scheduling

| Container Size | Plugs/Pot | Crop Time | Season | PGR |
|------------------------------|-----------|---------------|--------|--|
| 5"/6"/1 Gallon/15 cm | 1 (ppp) | 11-13 (weeks) | Spring | chlormequat chloride 300-500 ppm Spray |
| 12" Pot or HB/5 Gallon/30 cm | 4 (ppp) | 14-15 (weeks) | Spring | chlormequat chloride 300-500 ppm Spray |

Fertiliser Notation

Moderate fertilization levels are required. It is advisable to fertilize several times with low concentrations weekly. Fertilize crop, alternating with 150 to 200 ppm nitrogen, using a complete and potassium balanced fertilizer. Keep ammonium levels low to avoid damaging the roots. At high nitrogen levels, the foliage can become very big. Avoid pH above 6.0, as high pH causes iron deficiency. Apply chelated iron if chlorosis becomes a problem. To prevent magnesium deficiency, apply magnesium sulphate 1 to 2 times. Additional foliage fertilization with potassium supports compact plant growth, and the foliage will be a dark green color.

Common Problems


Monitor Aphids and thrips; preventative programs for powdery mildew are beneficial. Monitor Botrytis.

Finishing Key Tips


Do not overwater plants to prevent root issues. Keep humidity low (between 40 to 60%) to avoid problems with powdery mildew. For height control, a negative DIF of 2 to 3 degrees is very effective to prevent stretching of plants. When using a negative DIF, less to no PGRs are needed.

NOTE: Growers should use the information presented here as guidelines only. PanAmerican Seed recommends that growers conduct a trial of products under their own conditions. Crop times will vary depending on the climate, location, time of year, and greenhouse environmental conditions. It is the responsibility of the grower to confirm the treatment is available in their region as well as read and follow all the current label directions relating to the products. Nothing herein shall be deemed a warranty or guaranty by PanAmerican Seed of any products listed herein. PanAmerican Seed's terms and conditions of sale shall apply to all products listed herein.

Variety Pictures



Lace Pink



Lace Red



PanAmerican Seed Co.
622 Town Road, West Chicago, Illinois, USA, 60185
+1 800-231-7065 PanAmSeed.com

™ denotes a trademark of and ® denotes a registered trademark of Ball Horticultural Company in the US. It may also be registered in other countries.
©2025 Ball Horticultural Company