

## Fuseables® Bacopa Series

(*Sutera cordata*)

Annuals Culture (revised 08/14/19)

### Supplied as Fuseables® Precision™ Multi-Pellets.

These easy-to-produce, two to three-variety mixes are ideal for colour bowls, hanging baskets and large patio containers. Precision™ Multi-Pellets make it easy to grow beautiful combos – a “garden in a pellet!” We recommend 1 to 3 plants per 10-in./25-cm container and 3 to 5 plants per 12-in./30-cm container.

## General Information

Exposure	Bloom Season	Height	Spread	Spacing
Sun	Early Spring, Spring, Late Spring, Summer, Late Summer, Autumn	2-6 in. (5-15 cm)	12-18 in. (30-46 cm)	8-20 in. (20-51 cm)

## Germination

Seed Form	Recommended Plug Size	Seeds/Cell	Plug Crop Weeks	Days to Germinate	Initial Media pH/EC (1:2)	Cover Seed
PMPL	128 288	1	4-5	4	5.5-6.0 pH 0.75 mmhos/cm	No

## Plug Production

	Stage 1	Stage 2	Stage 3	Stage 4
<b>Moisture</b>	Level 4	Level 3-4	Level 3-4	Level 3
<b>Temperature</b>	68-73°F (20-23°C)	68-70°F (20-21°C)	65-70°F (18-21°C)	59-65°F (15-18°C)
<b>Light</b>	Light	1,000-2,500 f.c. (10,800-26,900 Lux)	1,000-2,500 f.c. (10,800-26,900 Lux)	2,500-5,000 f.c. (26,900-53,800 Lux)
<b>Fertilizer</b>		Less than 100 ppm N (Less than 0.7 EC)	100 to 175 ppm N (0.7 to 1.2 EC)	100 to 175 ppm N (0.7 to 1.2 EC)

## Growing on to Finish

Growing on Temperature	Target Media pH/EC (1:2)	Fertilizer	Daylength
(day) 59-76°F (15-24°C) (night) 55-60°F (13-16°C)	5.8-6.2 pH 1.2-1.5 mmhos/cm	175 to 225 ppm N (1.2 to 1.5 EC)	Day Neutral

## Crop Scheduling

Container Size	Plugs/Pot	Crop Time	Season	PGR
5"/6"/1 Gallon	1 (ppp)	6-7 (weeks)	Spring	daminozide 1,000-1,500 ppm Spray
8"/2 Gallon	3 (ppp)	6-7 (weeks)	Spring	daminozide 1,000-1,500 ppm Spray
10" Pot or HB/3 Gallon	4 (ppp)	6-7 (weeks)	Spring	daminozide 1,000-1,500 ppm Spray
12" Pot or HB/5 Gallon	4 (ppp)	6-7 (weeks)	Spring	daminozide 1,000-1,500 ppm Spray

## Fertilizer Notation

Fertilize with a predominately nitrate-based fertilizer with low phosphorus.

## Common Problems

Insects: Check for Whiteflies

## Finishing Key Tips

Avoid excessive watering and drought. Do not let plants wilt, as this will result in bud drop.

NOTE: Growers should use the information presented here as guidelines only. PanAmerican Seed recommends that growers conduct a trial of products under their own conditions. Crop times will vary depending on the climate, location, time of year, and greenhouse environmental conditions. It is the responsibility of the grower to confirm the treatment is available in their region as well as read and follow all the current label directions relating to the products. Nothing herein shall be deemed a warranty or guaranty by PanAmerican Seed of any products listed herein. PanAmerican Seed's terms and conditions of sale shall apply to all products listed herein.

## Variety Pictures

---



Utopia

PanAmericanSeed™

PanAmerican Seed Co.  
622 Town Road, West Chicago, Illinois, USA, 60185-2698  
630 231-1400 Fax: 630 231-3609 PanAmSeed.com

™ denotes a trademark of and © denotes a registered trademark of Ball Horticultural Company in the US. It may also be registered in other countries.  
©2021 Ball Horticultural Company