

Begonia Gryphon

(*Begonia x hybrida*)

Annuals Culture (revised 08/07/19)

Plug crop time: 8 to 9 weeks

Transplant to finish: 5 to 6 weeks

- Easy-care, low-water plant blends beauty with durability to make dramatic mixed or solo containers.
- Fits into existing begonia plug culture; easier and quicker to produce than vegetative foliage types.
- More tolerant of stress than Rex begonias for less shrink.
- Retailers appreciate the longer sales window and versatility for outdoor container and indoor potted plant sales.
- Lets consumers extend their enjoyment by bringing plants indoors at the end of the season.

General Information

Exposure	Bloom Season	Height	Spread	Spacing
Shade, Partial Sun	-	14-16 in. (36-41 cm)	16-18 in. (41-46 cm)	16-18 in. (41-46 cm)

Germination

Seed Form	Recommended Plug Size	Seeds/Cell	Plug Crop Weeks	Days from 50% to maximum germination	Initial Media pH/EC (1:2)	Cover Seed
MPL	288	1	8-9	10-12	5.8-6.2 pH 0.5 mmhos/cm	No

Plug Production

	Stage 1	Stage 2	Stage 3	Stage 4
Moisture	Level 5	Level 4-5	Level 3-4	Level 3
Temperature	72-78°F (22-26°C)	71-76°F (22-24°C)	68-73°F (20-23°C)	65-67°F (18-19°C)
Light	Light	2,000-2,500 f.c. (21,500-26,900 Lux)	2,500-5,000 f.c. (26,900-53,800 Lux)	2,500-5,000 f.c. (26,900-53,800 Lux)
Fertiliser		Less than 100 ppm N (Less than 0.7 EC)	100 to 175 ppm N (0.7 to 1.2 EC)	100 to 175 ppm N (0.7 to 1.2 EC)

Propagation Key Tips

A saturated media and high relative humidity are critical to successful germination.

Growing on to Finish

Growing on Temperature	Target Media pH/EC (1:2)	Fertiliser	Daylength
(day) 65-75°F (18-24°C) (night) 62-67°F (17-19°C)	5.4-6.0 pH 1.0 mmhos/cm	175 to 225 ppm N (1.2 to 1.5 EC)	Facultative Short Day

Daylength Notation

Gryphon is a foliage plant, but plant could flower when grown under a daylength of 11 hours or shorter. Under daylength longer than 11 hours, flowering will be significantly delayed or plants will never flower.

Crop Scheduling

Container Size	Plugs/Pot	Crop Time	Season	PGR
4"/4.5"/Quart/10 cm	1 (ppp)	5-6 (weeks)	Summer	-
5"/6"/1 Gallon/15 cm	2-3 (ppp)	5-6 (weeks)	Summer	-
8"/2 Gallon/20 cm	3 (ppp)	7-8 (weeks)	Summer	-
12" Pot or HB/5 Gallon/30 cm	3-4 (ppp)	9-11 (weeks)	Summer	-

Chemical Sensitivity

Avoid using Cycocel alone, as it can cause phytotoxicity. Also use caution with Bonzi, Topflor and Sumagic, as they can stunt plants.

Finishing Key Tips

To avoid flowering as a foliage plant, grow under daylength longer than 11 hours.

NOTE: Growers should use the information presented here as guidelines only. PanAmerican Seed recommends that growers conduct a trial of products under their own conditions. Crop times will vary depending on the climate, location, time of year, and greenhouse environmental conditions. It is the responsibility of the grower to confirm the treatment is available in their region as well as read and follow all the current label directions relating to the products. Nothing herein shall be deemed a warranty or guaranty by PanAmerican Seed of any products listed herein. PanAmerican Seed's terms and conditions of sale shall apply to all products listed herein.

Variety Pictures



Gryphon



Gryphon



Gryphon



PanAmerican Seed Co.
622 Town Road, West Chicago, Illinois, USA, 60185
+1 800-231-7065 PanAmSeed.com

™ denotes a trademark of and ® denotes a registered trademark of Ball Horticultural Company in the US. It may also be registered in other countries.
©2024 Ball Horticultural Company