

## Dahlia Figaro™ Series

(*Dahlia x hybrida*)

Annuals Culture (revised 01/12/21)

**Plug crop time:** 4 to 5 weeks

**Transplant to finish:** 5 to 6 weeks

- Premium pack item is ideal for borders and containers.
- Each variety contains several shades of that colour.

### General Information

Exposure	Bloom Season	Height	Spread	Spacing
Partial Sun	Spring, Late Spring, Summer, Autumn	14-18 in. (36-46 cm)	10-12 in. (25-30 cm)	10-14 in. (25-36 cm)

### Germination

Seed Form	Recommended Plug Size	Seeds/Cell	Plug Crop Weeks	Days from 50% to maximum germination	Initial Media pH/EC (1:2)	Cover Seed
RAW	288	1	4-5	3-7	5.5-5.8 pH 0.5-0.75 mmhos/cm	Yes

### Plug Production

	Stage 1	Stage 2	Stage 3	Stage 4
<b>Moisture</b>	Level 3	Level 3	Level 2-3	Level 2-3
<b>Temperature</b>	68-73°F (20-23°C)	68-73°F (20-23°C)	65-70°F (18-21°C)	65-70°F (18-21°C)
<b>Light</b>		1,500-3,000 f.c. (16,100-32,300 Lux)	1,500-3,000 f.c. (16,100-32,300 Lux)	2,500-4,000 f.c. (26,900-43,100 Lux)
<b>Fertilizer</b>		Less than 100 ppm N (Less than 0.7 EC)	100 to 175 ppm N (0.7 to 1.2 EC)	100 to 175 ppm N (0.7 to 1.2 EC)

### Propagation Key Tips

If a germ chamber is used, move trays to the greenhouse at first sign of germination.

### Growing on to Finish

Growing on Temperature	Target Media pH/EC (1:2)	Fertilizer	Daylength
(day) 52-60°F (11-16°C) (night) 52-60°F (11-16°C)	5.8-6.2 pH 0.75-1.0 mmhos/cm	225 to 300 ppm N (1.5 to 2.0 EC)	Obligate Long Day

### Daylength Notation

Figaro dahlias need to grow under less than 14 hours daylength; growing in 14 hours or more will result in issues with flower initiation.

### Crop Scheduling

Container Size	Plugs/Pot	Crop Time	Season	PGR
306 Pack/1801	1 (ppp)	5-6 (weeks)	Spring	-
4"/4.5"/Quart/10 cm	1 (ppp)	5-6 (weeks)	Spring	-

## Chemical Sensitivity

A-Rest, Bonzi, B-Nine/Alar and Cycocel can be used for plant control.


## Finishing Key Tips

Very responsive to B-Nine/Alar. Also responsive to day/night temperature differential (DIF), and plants are shorter with a negative DIF.

NOTE: Growers should use the information presented here as guidelines only. PanAmerican Seed recommends that growers conduct a trial of products under their own conditions. Crop times will vary depending on the climate, location, time of year, and greenhouse environmental conditions. It is the responsibility of the grower to confirm the treatment is available in their region as well as read and follow all the current label directions relating to the products. Nothing herein shall be deemed a warranty or guaranty by PanAmerican Seed of any products listed herein. PanAmerican Seed's terms and conditions of sale shall apply to all products listed herein.

## Variety Pictures

---



Orange Shades



Red Shades



Violet Shades



White



Yellow Shades



Mixture

PanAmericanSeed™

PanAmerican Seed Co.  
622 Town Road, West Chicago, Illinois, USA, 60185-2698  
630 231-1400 Fax: 630 231-3609 PanAmSeed.com

™ denotes a trademark of and ® denotes a registered trademark of Ball Horticultural Company in the US. It may also be registered in other countries.  
©2021 Ball Horticultural Company