

Pansy Frizzle Sizzle F₁ Series

(*Viola x wittrockiana*)

Annuals Culture (revised 02/25/19)

Plug crop time: 5 weeks

Transplant to finish: Spring 6 to 8 weeks, Autumn 4 to 6 weeks

- PanAmerican Seed is the only professional supplier of this unique, frilly flower form – count on reliable seed supply.
- Great for cool-season Winter and early Spring end cap displays.
- Looks great in containers and “close-up” landscapes.
- Stronger ruffling occurs in cool conditions; Spring production is recommended.

General Information

Exposure	Bloom Season	Height	Spread	Spacing
Partial Sun, Sun	Early Spring, Spring, Autumn	6-8 in. (15-20 cm)	8-10 in. (20-25 cm)	6-8 in. (15-20 cm)

Germination

Seed Form	Recommended Plug Size	Seeds/Cell	Plug Crop Weeks	Days to Germinate	Initial Media pH/EC	Cover Seed
PRM, RAW	288	1	5	3-4	5.5-5.8 pH 0.75 mmhos/cm	Yes

Plug Production

	Stage 1	Stage 2	Stage 3	Stage 4
Moisture	Level 4	Level 3-4	Level 3-4	Level 2-4
Temperature	65-70°F (18-21° C)	60-65°F (16-18°C)	60-65°F (16-18°C)	55-60°F (13-16°C)
Light	Optional	2,500 f.c. (26,900 Lux)	2,500 f.c. (26,900 Lux)	5,000 f.c. (53,800 Lux)
Fertilizer		Less than 100 ppm N - Less than 0.7 EC	100 to 175 ppm N - 0.7 to 1.2 EC	100 to 175 ppm N - 0.7 to 1.2 EC
PGR			ancymidol/5-10 ppm/Spray daminozide/1,500-2,500 ppm/Spray	ancymidol/5-10 ppm/Spray daminozide/1,500-2,500 ppm/Spray

Fertilizer Notation

It is best to use a nitrate-based fertilizer with low phosphorus to reduce stretch. Maintain a media pH of 5.5 to 5.8 and EC at 0.7 to 1.0 mS/cm (1:2 extraction). A higher pH (greater than 6.2) can induce Boron deficiency and also encourages fungal black root rot caused by *Thielaviopsis* sp.

Propagation Key Tips

Adjust PGR rates and frequency of application depending on local conditions in Stages 3 and 4.

Growing on to Finish

Growing on Temperature	Target Media pH/EC	Fertilizer	Daylength
(day) 60-70°F (16-21°C) (night) 50-55°F (10-13°C)	5.5-5.8 pH 1.5-2.0 mmhos/cm	175 to 225 ppm N - 1.2 to 1.5 EC	Facultative Long Day

Crop Scheduling

Container Size	Plugs/Pot	Crop Time	Season	PGR
5"/6"/1 Gallon	3 (ppp)	6-8 (weeks)	Early Spring	daminozide/chlormequat chloride tank mix 1,500/150-2,500/250 ppm Spray
10" Pot or HB/3 Gallon	7-9 (ppp)	6-8 (weeks)	Early Spring	daminozide/chlormequat chloride tank mix 1,500/150-2,500/250 ppm Spray
5"/6"/1 Gallon	3 (ppp)	4-6 (weeks)	Autumn	daminozide/chlormequat chloride tank mix 2,500/250-5,000/500 ppm Spray
10" Pot or HB/3 Gallon	7-9 (ppp)	4-6 (weeks)	Autumn	daminozide/chlormequat chloride tank mix 2,500/250-5,000/500 ppm Spray

Common Problems

Insects: Fungus Gnats and Shore Flies in plug stage and Aphids in early stages after transplant. Disease: Damping off, Black Root Rot, Foliar Leaf Spots, Botrytis

Finishing Key Tips


Adjust PGR rates and frequency of application depending on local conditions. If growing frost-free (northern overwinter culture from Autumn transplant) plan for crop times of 16 to 18 weeks from transplant to finish. For maximum ruffled edge, schedule for late Autumn, Winter and Spring programs. Heat reduces the ruffled edge and colour contrast.

NOTE: Growers should use the information presented here as guidelines only. PanAmerican Seed recommends that growers conduct a trial of products under their own conditions. Crop times will vary depending on the climate, location, time of year, and greenhouse environmental conditions. It is the responsibility of the grower to confirm the treatment is available in their region as well as read and follow all the current label directions relating to the products. Nothing herein shall be deemed a warranty or guaranty by PanAmerican Seed of any products listed herein. PanAmerican Seed's terms and conditions of sale shall apply to all products listed herein.


Variety Pictures



Blue



Burgundy



Lemonade



Lemonberry



Orange



Raspberry



Yellow



Yellow-Blue Swirl



Mixture

PanAmericanSeed™

PanAmerican Seed Co.
622 Town Road, West Chicago, Illinois, USA, 60185-2698
630 231-1400 Fax: 630 231-3609 PanAmSeed.com

™ denotes a trademark of and ® denotes a registered trademark of Ball Horticultural Company in the US. It may also be registered in other countries.
©2019 Ball Horticultural Company