GrowerFacts

PanAmerican Seed®

Pansy Panola® XP F_1 Series

(Viola x wittrockiana)

Annuals Culture (revised 10/26/23)

Plug crop time: 5 weeks
Transplant to finish: Spring 4 to 5 weeks, Autumn 3 to 4 weeks

- It features a tight 5 to 7-day flowering window, very uniform habit and shorter peduncles across a broad array of colours.
- These colours outperform standard medium-flowered varieties in all kinds of conditions.
- Exceptional overwintering ability makes Panola XP the best choice for overwintering guarantee pansy programs.
- Free-flowering, well-branched and compact, the early-blooming plants maintain their garden height with less stretch in warm climates.
- Plants produce an abundance of blooms for good flower show in the landscape.
- Landscaper-friendly clear colours are perfect for mixing custom blends.
- Ideal for packs and 4-in. (9 to 10.5-cm) pots.

General Information

Exposure	Bloom Season	Height	Spread	Spacing
Partial Sun	Early Spring, Spring, Autumn	6-8 in. (15-20 cm)	8-10 in. (20-25 cm)	6-8 in. (15-20 cm)

Germination

Seed Form	Recommended Plug Size	Seeds/Cell	Plug Crop Weeks	Days from 50% to maximum germination	Initial Media pH/EC (1:2)	Cover Seed
PRM, RAW	288	1	5	3-4	5.5-5.8 pH 0.75 mmhos/cm	Yes

Plug Production

	Stage 1	Stage 2	Stage 3	Stage 4
Moisture	Level 4	Level 3-4	Level 3-4	Level 2-4
Temperature	65-70°F (18-21°C)	60-65°F (16-18°C)	60-65°F (16-18°C)	55-60°F (13-16°C)
Light	Optional	2,500 f.c. (26,900 Lux)	2,500 f.c. (26,900 Lux)	5,000 f.c. (53,800 Lux)
Fertiliser		Less than 100 ppm N (Less than 0.7 EC)	100 to 175 ppm N (0.7 to 1.2 EC)	100 to 175 ppm N (0.7 to 1.2 EC)
PGR			ancymidol/5-10 ppm/Spray daminozide/1,500-2,500 ppm/Spray	ancymidol/5-10 ppm/Spray daminozide/1,500-2,500 ppm/Spray

Fertiliser Notation

It is best to use a nitrate-based fertilizer with low phosphorus to reduce stretch. Maintain a media pH of 5.5 to 5.8 and EC at 0.7 to 1.0 mS/cm (1:2 extraction). A higher pH (greater than 6.2) can induce Boron deficiency and also encourages fungal black root rot caused by Thielaviopsis sp.

Propagation Key Tips

Adjust PGR rates and frequency of application depending on local conditions in Stages 3 and 4.

Growing on to Finish

Growing on Temperature	Target Media pH/EC (1:2)	Fertiliser	Daylength
(day) 60-70°F (16-21°C) **(night)** 50-55°F (10-13°C)	5.5-5.8 pH 1.5-2.0 mmhos/cm	100 to 175 ppm N (0.7 to 1.2 EC)	Facultative Long Day

Crop Scheduling

Container Size	Plugs/Pot	Crop Time	Season	PGR
Cell Pack	1 (ppp)	5-6 (weeks)	Early Spring	daminozide/chlormequat chloride tank mix 1,500/150-2,500/250 ppm Spray
4"/4.5"/Quart/10 cm	1 (ppp)	5-6 (weeks)	Early Spring	daminozide/chlormequat chloride tank mix 1,500/150-2,500/250 ppm Spray
Cell Pack	1 (ppp)	3-4 (weeks)	Autumn	daminozide/chlormequat chloride tank mix 2,500/500-5,000/500 ppm Spray
Cell Pack	1 (ppp)	3-4 (weeks)	Autumn	paclobutrazol 0.1-0.125 ppm Drench
4"/4.5"/Quart/10 cm	1 (ppp)	4-5 (weeks)	Autumn	daminozide/chlormequat chloride tank mix 2,500/500-5,000/500 ppm Spray
4"/4.5"/Quart/10 cm	1 (ppp)	3-4 (weeks)	Autumn	paclobutrazol 0.1-0.125 ppm Drench

Common Problems

Insects: Fungus Gnats and Shore Flies in plug stage and Aphids in early stages after transplant. Disease: Damping off, Black Root Rot, Foliar Leaf Spots, Botrytis

Finishing Key Tips

Adjust PGR rates and frequency of application depending on local conditions. If growing frost-free (northern overwinter culture from Autumn transplant), plan for crop times of 16 to 18 weeks from transplant to finish.

NOTE: Growers should use the information presented here as guidelines only. PanAmerican Seed recommends that growers conduct a trial of products under their own conditions. Crop times will vary depending on the climate, location, time of year, and greenhouse environmental conditions. It is the responsibility of the grower to confirm the treatment is available in their region as well as read and follow all the current label directions relating to the products. Nothing herein shall be deemed a warranty or guaranty by PanAmerican Seed of any products listed herein. PanAmerican Seed's terms and conditions of sale shall apply to all products listed herein.

Variety Pictures

Beaconsfield

Deep Blue Blotch

Deep Orange

Purple

Rose Picotee

Scarlet

True Blue

White

Yellow

Yellow Blotch

Citrus Mixture

Clear Mixture

Mixture

PanAmerican Seed Co.
622 Town Road,West Chicago, Illinois, USA, 60185
+1 800-231-7065 PanAmSeed.com

™ denotes a trademark of and ® denotes a registered trademark of Ball Horticultural Company in the US. It may also be registered in other countries.
©2024 Ball Horticultural Company