

Petunia Debonair™ F₁ Series

(*Petunia x hybrida*)

Annuals Culture (revised 02/24/20)

The designer Debonair and Sophistica collections combine beautiful flowers in special one-of-a-kind colours with professional seed quality. These premium petunias can command a higher price point at retail and open the door to decorator opportunities.

Plug crop time: 4 to 5 weeks

Transplant to finish: 5 to 6 weeks

- Plants are well-matched for timing for easier scheduling and shipping.
- Mounded, upright habit is ideal for premium pots and larger containers.

General Information

Exposure	Bloom Season	Height	Spread	Spacing
Sun	Spring, Late Spring, Summer	10-15 in. (25-38 cm)	10-12 in. (25-30 cm)	10-14 in. (25-36 cm)

Germination

Seed Form	Recommended Plug Size	Seeds/Cell	Plug Crop Weeks	Days to Germinate	Initial Media pH/EC	Cover Seed
PEL, RAW	288	1	4-5	3-5	5.5-5.8 pH 0.75-1.0 mmhos/cm	No

Plug Production

	Stage 1	Stage 2	Stage 3	Stage 4
Moisture	Level 4-5	Level 3-4	Level 2-4	Level 2-4
Temperature	72-76°F (22-24°C)	68-75°F (20-24°C)	65-70°F (18-21°C)	60-65°F (16-18°C)
Light	Light	1,000-2,500 f.c. (10,800-26,900 Lux)	2,500-4,000 f.c. (26,900-43,100 Lux)	2,500-5,000 f.c. (26,900-53,800 Lux)
Fertilizer		Less than 100 ppm N - Less than 0.7 EC	100 to 175 ppm N - 0.7 to 1.2 EC	100 to 175 ppm N - 0.7 to 1.2 EC
PGR			daminozide/2,500-5,000 ppm/Spray	daminozide/2,500-5,000 ppm/Spray

Propagation Key Tips

PGR options include paclobutrazol or flurprimidol.

Growing on to Finish

Growing on Temperature	Target Media pH/EC	Fertilizer	Daylength
(day) 62-70°F (17-21°C) (night) 55-65°F (13-18°C)	5.5-6.2 pH 1.0-1.5 mmhos/cm	100 to 175 ppm N - 0.7 to 1.2 EC	Facultative Long Day

Daylength Notation

All varieties can flower successfully at 10-hour daylengths.

Crop Scheduling

Container Size	Plugs/Pot	Crop Time	Season	PGR
4"/4.5"/Quart	1 (ppp)	5-6 (weeks)	Spring	paclobutrazol 2-3 ppm Drench
10" Pot or HB/3 Gallon	3 (ppp)	6-7 (weeks)	Spring	paclobutrazol 2-3 ppm Drench
4"/4.5"/Quart	1 (ppp)	3-5 (weeks)	Summer	paclobutrazol 2-3 ppm Drench

Chemical Sensitivity

Avoid using daminozide on Black Cherry, as this may impact flower colour.

Common Problems


Insects: Thrips, Aphids, Fungus Gnats, Shore Flies Diseases: Botrytis, Rhizoctonia, Pythium

Finishing Key Tips


Avoid using daminozide on Black Cherry, as this may impact flower colour. Options include paclobutrazol or flurprimidol.

NOTE: Growers should use the information presented here as guidelines only. PanAmerican Seed recommends that growers conduct a trial of products under their own conditions. Crop times will vary depending on the climate, location, time of year, and greenhouse environmental conditions. It is the responsibility of the grower to confirm the treatment is available in their region as well as read and follow all the current label directions relating to the products. Nothing herein shall be deemed a warranty or guaranty by PanAmerican Seed of any products listed herein. PanAmerican Seed's terms and conditions of sale shall apply to all products listed herein.


Variety Pictures



Black Cherry



Dusty Rose



Lime Green

PanAmericanSeed™

PanAmerican Seed Co.
622 Town Road, West Chicago, Illinois, USA, 60185-2698
630 231-1400 Fax: 630 231-3609 PanAmSeed.com

™ denotes a trademark of and © denotes a registered trademark of Ball Horticultural Company in the US. It may also be registered in other countries.
©2020 Ball Horticultural Company