

Phlox Promise Series

(*Phlox drummondii*)

Annuals Culture (revised 08/29/19)

Plug crop time: 4 to 5 weeks

Transplant to finish: 7 to 8 weeks

- Very floriferous plants feature double and semi-double flowers held well above the foliage.
- Strong outdoor vigour assures long-lasting garden performance.
- Well-suited to large pack and pot production.

General Information

Exposure	Bloom Season	Height	Spread	Spacing
Sun	Early Spring, Spring, Late Spring, Summer, Late Summer, Autumn	8 in. (20 cm)	8 in. (20 cm)	6-8 in. (15-20 cm)

Germination

Seed Form	Recommended Plug Size	Seeds/Cell	Plug Crop Weeks	Days to Germinate	Initial Media pH/EC	Cover Seed
RAW	288	1	4-5	3-5	5.8-6.0 pH 0.75 mmhos/cm	Yes

Plug Production

	Stage 1	Stage 2	Stage 3	Stage 4
Moisture	Level 4	Level 3-4	Level 3	Level 2-3
Temperature	65-70°F (18-21°C)	65-72°F (18-22°C)	60-65°F (16-18°C)	60-65°F (16-18°C)
Light	Dark	1,000 f.c. (10,800 Lux)	2,500-3,000 f.c. (26,900-32,300 Lux)	2,500-3,000 f.c. (26,900-32,300 Lux)
Fertilizer	Less than 100 ppm N - Less than 0.7 EC		100 to 175 ppm N - 0.7 to 1.2 EC	100 to 175 ppm N - 0.7 to 1.2 EC
PGR			daminozide/2,500 ppm/Spray	daminozide/2,500 ppm/Spray

Propagation Key Tips

Cover thoroughly with coarse vermiculite. Darkness is required for germination.

Growing on to Finish

Growing on Temperature	Target Media pH/EC	Fertilizer	Daylength
(day) 60-68°F (16-20°C) (night) 55-62°F (13-17°C)	5.5-6.2 pH 1.0-1.5 mmhos/cm	175 to 225 ppm N - 1.2 to 1.5 EC	Day Neutral

Crop Scheduling

Container Size	Plugs/Pot	Crop Time	Season	PGR
Cell Pack	1 (ppp)	6-7 (weeks)	Spring	-
4"/4.5"/Quart	1 (ppp)	7-8 (weeks)	Spring	-

Finishing Key Tips


If necessary, paclobutrazol 7.5 to 20 ppm Spray can be applied. Application rate and frequency of applications depends upon growth rate. Daminozide 2,500 ppm Spray is also effective.

NOTE: Growers should use the information presented here as guidelines only. PanAmerican Seed recommends that growers conduct a trial of products under their own conditions. Crop times will vary depending on the climate, location, time of year, and greenhouse environmental conditions. It is the responsibility of the grower to confirm the treatment is available in their region as well as read and follow all the current label directions relating to the products. Nothing herein shall be deemed a warranty or guaranty by PanAmerican Seed of any products listed herein. PanAmerican Seed's terms and conditions of sale shall apply to all products listed herein.


Variety Pictures




Lilac




Peach




Pink




Rose



Scarlet



White



Mixture

PanAmericanSeed™

PanAmerican Seed Co.
622 Town Road, West Chicago, Illinois, USA, 60185-2698
630 231-1400 Fax: 630 231-3609 PanAmSeed.com

™ denotes a trademark of and © denotes a registered trademark of Ball Horticultural Company in the US. It may also be registered in other countries.
©2020 Ball Horticultural Company