

## Phlox Ethnie Series

(*Phlox drummondii*)

Annuals Culture (revised 01/12/21)

**Plug crop time:** 4 to 5 weeks

**Transplant to finish:** 6 to 7 weeks

- Genetically compact series is an excellent choice for pot and pack programs, delivering uniform habit and earliness.
- Crop time is just 10 to 12 weeks.
- No PGRs are needed in production.

### General Information

Exposure	Bloom Season	Height	Spread	Spacing
Sun	Early Spring, Spring, Late Spring, Summer, Late Summer, Autumn	8 in. (20 cm)	8 in. (20 cm)	6-8 in. (15-20 cm)

### Germination

Seed Form	Recommended Plug Size	Seeds/Cell	Plug Crop Weeks	Days to Germinate	Initial Media pH/EC (1:2)	Cover Seed
RAW	288	1	4-5	3-5	5.8-6.0 pH 0.75 mmhos/cm	Yes

### Plug Production

	Stage 1	Stage 2	Stage 3	Stage 4
<b>Moisture</b>	Level 4	Level 3-4	Level 3	Level 2-3
<b>Temperature</b>	65-70°F (18-21°C)	65-72°F (18-22°C)	60-65°F (16-18°C)	60-65°F (16-18°C)
<b>Light</b>	Dark	1,000 f.c. (10,800 Lux)	2,500-3,000 f.c. (26,900-32,300 Lux)	2,500-3,000 f.c. (26,900-32,300 Lux)
<b>Fertilizer</b>		Less than 100 ppm N (Less than 0.7 EC)	100 to 175 ppm N (0.7 to 1.2 EC)	100 to 175 ppm N (0.7 to 1.2 EC)
<b>PGR</b>			daminozide/2,500 ppm/Spray	daminozide/2,500 ppm/Spray

### Propagation Key Tips

Cover thoroughly with coarse vermiculite. Darkness is required for germination.

### Growing on to Finish

Growing on Temperature	Target Media pH/EC (1:2)	Fertilizer	Daylength
<b>(day)</b> 60-68°F (16-20°C) <b>(night)</b> 55-62°F (13-17°C)	5.5-6.2 pH 1.0-1.5 mmhos/cm	175 to 225 ppm N (1.2 to 1.5 EC)	Day Neutral

### Crop Scheduling

Container Size	Plugs/Pot	Crop Time	Season	PGR
Cell Pack	1 (ppp)	5-6 (weeks)	Spring	-
4"/4.5"/Quart	1 (ppp)	6-7 (weeks)	Spring	-

## Finishing Key Tips

If necessary, paclobutrazol 7.5 to 20 ppm Spray can be applied. Application rate and frequency of applications depends upon growth rate. Daminozide 2,500 ppm Spray is also effective.

NOTE: Growers should use the information presented here as guidelines only. PanAmerican Seed recommends that growers conduct a trial of products under their own conditions. Crop times will vary depending on the climate, location, time of year, and greenhouse environmental conditions. It is the responsibility of the grower to confirm the treatment is available in their region as well as read and follow all the current label directions relating to the products. Nothing herein shall be deemed a warranty or guaranty by PanAmerican Seed of any products listed herein. PanAmerican Seed's terms and conditions of sale shall apply to all products listed herein.

## Variety Pictures

---



Dark Rose



Light Blue



Purple Violet



Salmon



Scarlet



White



Mixture

PanAmericanSeed™

PanAmerican Seed Co.  
622 Town Road, West Chicago, Illinois, USA, 60185-2698  
630 231-1400 Fax: 630 231-3609 [PanAmSeed.com](http://PanAmSeed.com)

™ denotes a trademark of and © denotes a registered trademark of Ball Horticultural Company in the US. It may also be registered in other countries.  
©2021 Ball Horticultural Company