GrowerFacts

ColorGrass® Carex Red Rooster

(Carex buchananii)

Plug crop time: 6 to 7 weeks Transplant to finish: 8 to 9 weeks

Red-bronze grass has curled tips and upright habit.

Annuals Culture (revised 01/14/21)

General Information

Exposure	Bloom Season	Height	Spread	Spacing
Sun	Summer, Late Summer	20-30 in. (51-76 cm)	12 in. (30 cm)	61-91 in. (155-231 cm)

Germination

Seed Form	Recommended Plug Size	Seeds/Cell	Plug Crop Weeks	Days from 50% to maximum germination	Initial Media pH/EC (1:2)	Cover Seed
MPL	288	1	6-7	7-10	5.8-6.2 pH 0.75 mmhos/cm	No

Plug Production

	Stage 1	Stage 2	Stage 3	Stage 4
Moisture	Level 4	Level 3	Level 2-3	Level 2-3
Temperature	74-79°F (23-26°C)	68-72°F (20-22°C)	68-72°F (20-22°C)	65-67°F (18-19°C)
Light	Optional	1,000-2,500 f.c. (10,800-26,900 Lux)	1,000-2,500 f.c. (10,800-26,900 Lux)	2,500-5,000 f.c. (26,900-53,800 Lux)
Fertiliser		Less than 100 ppm N (Less than 0.7 EC)	100 to 175 ppm N (0.7 to 1.2 EC)	100 to 175 ppm N (0.7 to 1.2 EC)

Fertiliser Notation

Use a nitrate-based fertilizer with low phosphorus.

Growing on to Finish

Growing on Temperature	Target Media pH/EC (1:2)	Fertiliser	Daylength
(day) 66-74°F (19-23°C)	5.5-6.2 pH	175 to 225 ppm N	Day Neutral
(night) 64-66°F (18-19°C)	1.2-1.5 mmhos/cm	(1.2 to 1.5 EC)	

Crop Scheduling

Container Size	Plugs/Pot	Crop Time	Season	PGR
306 Pack/1801	1 (ppp)	8-9 (weeks)	Spring	-
4"/4.5"/Quart/10 cm	1 (ppp)	8-9 (weeks)	Spring	-
5"/6"/1 Gallon/15 cm	3 (ppp)	9-10 (weeks)	Spring	-

Fertiliser Notation

Use a nitrate-based fertilizer with low phosphorus; avoid using excessive ammonia, nitrogen-based fertilizer and overfeeding, as these will result in less upright plants.

Common Problems

Insects: Aphids, especially after Winter regrowth Disease: Root Rot, when grown too wet

NOTE: Growers should use the information presented here as guidelines only. PanAmerican Seed recommends that growers conduct a trial of products under their own conditions. Crop times will vary depending on the climate, location, time of year, and greenhouse environmental conditions. It is the responsibility of the grower to confirm the treatment is available in their region as well as read and follow all the current label directions relating to the products. Nothing herein shall be deemed a warranty or guaranty by PanAmerican Seed of any products listed herein. PanAmerican Seed's terms and conditions of sale shall apply to all products listed herein.

Variety Pictures







Red Rooster

Red Rooster

Red Rooster