# Verbena Santos Purple

(Verbena rigida)

Perennials Culture (revised 10/31/23)

Plug crop time: 6 to 7 weeks Transplant to finish: 11 to 13 weeks

- One of the best tough and long-lasting landscape perennials available very heat and drought tolerant, plus exceptional performance in cool climates
- Compact and upright, well-branched plants bloom late Spring through Summer.
- High powdery mildew resistance.
- · Ideal for use in combo planters.

### **General Information**

Exposure	Bloom Season	Height	Spread	Spacing
Sun	Late Spring, Summer, Late Summer, Autumn	12 in. (30 cm)	48 in. (122 cm)	16 in. (41 cm)

## **Germination**

Seed Form	Recommended Plug Size	Seeds/Cell	Plug Crop Weeks	Days from 50% to maximum germination	Initial Media pH/EC (1:2)	Cover Seed
PRM	288 128*	4 4	6-7 8	7-10	5.5-6.2 pH 0.75 mmhos/cm	Yes

<sup>\*</sup> Recommend plug size for forcing.

## **Plug Production**

	Stage 1	Stage 2	Stage 3	Stage 4
Moisture	Level 4	Level 3	Level 2	Level 2
Temperature	68-72° F (20-22° C)	65-68°F (18-20°C)	60-65°F (16-18°C)	60-65°F (16-18°C)
Light	Dark	2,000-2,500 f.c. (21,500- 26,900 Lux)	2,000-2,500 f.c. (21,500-26,900 Lux)	4,000-5,000 f.c. (43,100-53,800 Lux)
Fertiliser		100 to 175 ppm N (0.7 to 1.2 EC)	100 to 175 ppm N (0.7 to 1.2 EC)	100 to 175 ppm N (0.7 to 1.2 EC)
PGR			daminozide/chlormequat chloride tank mix/1,500/200-2,500/300 ppm/Spray	daminozide/chlormequat chloride tank mix/2,500/300-3,750/500 ppm/Spray

#### Vernalisation

No

## **Propagation Key Tips**

For forcing info: see Perennials Forcing Guide. Spray damp-off fungicide.

## **Growing on to Finish**

Growing on Temperature	Target Media pH/EC (1:2)	Fertiliser	Daylength
(day) 65-68°F (18-20°C)	5.5-6.2 pH	175 to 225 ppm N	Facultative Long Day
(night) 60-65°F (16-18°C)	1.3-1.5 mmhos/cm	(1.2 to 1.5 EC)	

#### **Daylength Notation**

For forcing info: See Perennials Forcing Guide.

## **Crop Scheduling**

Crop Type	Container Size	Plugs/Pot	Crop Time	Season	PGR
Annual	4"/4.5"/Quart/10 cm	1 (ppp)	11-14 (weeks)	Late Spring	daminozide/chlormequat chloride tank mix 2,000-2,500 ppm Spray
Annual	4"/4.5"/Quart/10 cm	1 (ppp)	11-14 (weeks)	Late Spring	daminozide 2,000-2,500 ppm Spray
Annual	5"/6"/1 Gallon/15 cm	3-4 (ppp)	10-13 (weeks)	Summer	daminozide/chlormequat chloride tank mix 2,000-750 ppm Spray
Annual	5"/6"/1 Gallon/15 cm	3-4 (ppp)	10-13 (weeks)	Summer	daminozide 2,000-2,500 ppm Spray
Annual	5"/6"/1 Gallon/15 cm	3-4 (ppp)	11-14 (weeks)	Late Spring	daminozide/chlormequat chloride tank mix 2,000-750 ppm Spray
Annual	5"/6"/1 Gallon/15 cm	3-4 (ppp)	11-14 (weeks)	Late Spring	daminozide 2,000-2,500 ppm Spray

#### **Fertiliser Notation**

Apply fertiliser at rate 1 to 2 (100 to 125 ppm N/1.1 to 1.3 mmhos/cm EC) or 200 ppm as needed. Maintain the pH at 5.5 to 6.

#### **Common Problems**

Monitor for Powdery Mildew, Aphids and Thrips.

#### **Finishing Key Tips**

Grow dry and light. Relatively high fertilisation. Avoid high N. Prevent Mg and Fe deficiency.

NOTE: Growers should use the information presented here as guidelines only. PanAmerican Seed recommends that growers conduct a trial of products under their own conditions. Crop times will vary depending on the climate, location, time of year, and greenhouse environmental conditions. It is the responsibility of the grower to confirm the treatment is available in their region as well as read and follow all the current label directions relating to the products. Nothing herein shall be deemed a warranty or guaranty by PanAmerican Seed of any products listed herein. PanAmerican Seed's terms and conditions of sale shall apply to all products listed herein.

# **Variety Pictures**









Santos Purple

Santos Purple

Santos Purple