

## Gerbera Mega Revolution™ F<sub>1</sub> Series

(*Gerbera jamesonii*)

Potted Plants Culture (revised 01/25/19)

**Plug crop time:** 6 to 7 weeks

**Transplant to finish:** 8 to 10 weeks

- Produces manageable, larger-sized plants with very large flowers – beats competing series for colour range, bloom size and seed quality.
- Ideal for larger container programs.
- Supplied with industry-leading seed quality.

### General Information

| Exposure    | Bloom Season   | Height                 | Spread                  | Spacing                 |
|-------------|--|------------------------|-------------------------|-------------------------|
| Partial Sun | Early Spring, Spring, Late Spring, Summer, Late Summer, Autumn, Winter | 6-10 in.<br>(15-25 cm) | 12-14 in.<br>(30-36 cm) | 12-14 in.<br>(30-36 cm) |

### Germination

| Seed Form | Recommended Plug Size | Seeds/Cell | Plug Crop Weeks | Days to Germinate | Initial Media pH/EC            | Cover Seed  |
|-----------|-----------------------|------------|-----------------|-------------------|--------------------------------|-------------|
| COT       | 128                   | 1          | 6-7             | 4-7               | 5.5-5.8 pH<br>0.5-0.7 mmhos/cm | Light cover |

### Plug Production

|                    | Stage 1                                | Stage 2   | Stage 3   | Stage 4   |
|--------------------|--|---|---|---|
| <b>Moisture</b>    | Level 5                                | Level 4   | Level 3   | Level 3   |
| <b>Temperature</b> | 70-74°F<br>(21-23°C)                   | 68-70°F<br>(20-21°C)  | 68-70°F<br>(20-21°C)  | 66-68°F<br>(19-20°C)  |
| <b>Light</b>       | Optional                               | 6-8 mol·m <sup>-2</sup> ·d <sup>-1</sup><br>2,000-2,500 f.c.<br>(21,500-26,900 Lux) | 6-8 mol·m <sup>-2</sup> ·d <sup>-1</sup><br>2,000-2,500 f.c.<br>(21,500-26,900 Lux) | 6-8 mol·m <sup>-2</sup> ·d <sup>-1</sup><br>2,000-2,500 f.c.<br>(21,500-26,900 Lux) |
| <b>Fertilizer</b>  | Less than 100 ppm N - Less than 0.7 EC | Less than 100 ppm N - Less than 0.7 EC  | 100 to 175 ppm N - 0.7 to 1.2 EC  | 100 to 175 ppm N - 0.7 to 1.2 EC  |

### Propagation Key Tips

Don't water with cold water. Use water with a temperature at least 60°F/16°C or higher. Watering with cold water will result in deformed leaves and will disturb plant growth. Do not pot too deeply when transplanting, as this may result in Crown Rot. Space plants when the leaves of the plants are touching each other, generally 5 to 6 weeks after transplanting.

### Growing on to Finish

| Growing on Temperature   | Target Media pH/EC             | Fertilizer                       | Daylength             |
|--|--------------------------------|----------------------------------|-----------------------|
| <b>(day)</b> 66-68°F (19-20°C)<br><b>(night)</b> 62-66°F (17-19°C) | 5.5-6.0 pH<br>0.5-1.0 mmhos/cm | 100 to 175 ppm N - 0.7 to 1.2 EC | Facultative Short Day |

## Crop Scheduling

| Container Size | Plugs/Pot | Crop Time     | Season | PGR                              |
|----------------|-----------|---------------|--------|----------------------------------|
| 5"/6"/1 Gallon | 1 (ppp)   | 10-11 (weeks) | Spring | daminozide 1,000-2,500 ppm Spray |
| 5"/6"/1 Gallon | 1 (ppp)   | 8-9 (weeks)   | Summer | daminozide 1,000-2,500 ppm Spray |
| 5"/6"/1 Gallon | 1 (ppp)   | 10-11 (weeks) | Autumn | daminozide 1,000-2,500 ppm Spray |
| 5"/6"/1 Gallon | 1 (ppp)   | 11-12 (weeks) | Autumn | daminozide 1,000-2,500 ppm Spray |

### Fertilizer Notation

Reduce nitrogen (N) in fertilizer mix at bud initiation stage until flowering for better plant control.

### Common Problems

Gerbera is very sensitive to Thrips. Zero tolerance is recommended. Also watch for Whiteflies, Leaf Miners, Spider Mites and other insects. It is recommended to start spraying against insects right after transplant.

### Finishing Key Tips

Use 6-in. (15 cm) pots or larger. Fill pots all the way to a uniform soil level. When transplanting, do not plant too deep, as this may result in Crown Rot. Space plants when leaves are touching, generally 5 to 6 weeks after transplanting. To reduce stretching, use B-Nine/Alar (daminozide) at 1,000 to 2,500 ppm 1 to 2 times with an interval of 9 to 10 days. Stop use after seeing the first flower buds.

NOTE: Growers should use the information presented here as guidelines only. PanAmerican Seed recommends that growers conduct a trial of products under their own conditions. Crop times will vary depending on the climate, location, time of year, and greenhouse environmental conditions. It is the responsibility of the grower to confirm the treatment is available in their region as well as read and follow all the current label directions relating to the products. Nothing herein shall be deemed a warranty or guaranty by PanAmerican Seed of any products listed herein. PanAmerican Seed's terms and conditions of sale shall apply to all products listed herein.

## Variety Pictures



Bright Orange with Light Eye



Bright Red



Bright Rose with Dark Eye



Deep Rose with Light Eye



Orange with Dark Eye



Orange with Light Eye



Rose with Light Eye



Salmon Rose with Light Eye



Scarlet Red with Light Eye



Watermelon



White with Light Eye



Yellow with Dark Eye



Select Mixture

PanAmericanSeed™

PanAmerican Seed Co.  
622 Town Road, West Chicago, Illinois, USA, 60185-2698  
630 231-1400 Fax: 630 231-3609 [PanAmSeed.com](http://PanAmSeed.com)

™ denotes a trademark of and ® denotes a registered trademark of Ball Horticultural Company in the US. It may also be registered in other countries.  
©2019 Ball Horticultural Company