

Pepper (Edible Potted) Hot Tomato

(*Capsicum annuum*)

Vegetables Culture

- Unique assortment is bred for shorter production times.
- Compact growth means no PGRs are needed.
- Good retail shelf life.
- Excellent indoor/outdoor performance.

General Information

Exposure	Bloom Season	Height	Spread	Spacing
Sun	Summer	10-12 in. (25-30 cm)	8-10 in. (20-25 cm)	

Germination

Seed Form	Recommended Plug Size	Seeds/Cell	Plug Crop Weeks	Days to Germinate	Initial Media pH/EC	Cover Seed
RAW	288	1	3-4	5-6	5.4-5.8 pH 0.8-1.2 mmhos/cm	No

Plug Production

	Stage 1	Stage 2	Stage 3	Stage 4
Moisture	Level 4	Level 3-4	Level 2-4	Level 2-4
Temperature	70-73°F (21-23°C)	68-70°F (20-21°C)	68-70°F (20-21°C)	68-70°F (20-21°C)
Light	Optional	1,000-2,500 f.c. (10,800-26,900 Lux)	1,000-2,500 f.c. (10,800-26,900 Lux)	2,500-5,000 f.c. (26,900-53,800 Lux)
Fertilizer	Less than 100 ppm N - Less than 0.7 EC	100 to 175 ppm N - 0.7 to 1.2 EC	175 to 225 ppm N - 1.2 to 1.5 EC	175 to 225 ppm N - 1.2 to 1.5 EC
PGR			uniconazole/2.5 ppm/Spray	uniconazole/2.5 ppm/Spray

Propagation Key Tips

To ensure a uniform crop, growth regulator should be applied with a fine mist. For more vigorous varieties, an application of uniconazole 2.5 ppm at 10 to 12 days after sowing, repeated 2 weeks later if needed, will help control plant size. To bench germ: Make sure trays are watered before covering. Flat covering: first place one layer of reemay, then one layer of white plastic on top of that. Remove plastic after 8 to 10 days. Keep reemay wet. Remove reemay after another 2 to 3 days.

Growing on to Finish

Growing on Temperature	Target Media pH/EC	Fertilizer	Daylength
(day) 68-80°F (20-27°C) (night) 66-70°F (19-21°C)	5.5-5.9 pH 1.8-2.5 mmhos/cm	More than 300 ppm N - More than 2.0 EC	Day Neutral

Crop Scheduling

Container Size	Plugs/Pot	Crop Time	Season	PGR
5"/6"/1 Gallon	1 (ppp)	17 (weeks)	Spring	-
5"/6"/1 Gallon	1 (ppp)	15 (weeks)	Summer	-

Fertilizer Notation

Especially when plants are grown very dry, it's important to use higher EC approximately once a week when plants are watered from below. Ensure sufficient Ca by the end of the crop.

Common Problems

The most common pest with capsicum is Thrips. We recommend putting up sticky cards (1 per 20-ft./6-m table).

Finishing Key Tips

Plant plugs deeply. Water from below to keep surface of media dry. It is especially important to grow plants dry until flowering, allowing some wilt between watering. To optimize plant growth, space as soon as the plants touch. Keep the plants very dry from spacing to flowering to avoid stretch. Due to the weight of the fruit, we recommend staking plants to stabilize plants during shipment.

NOTE: Growers should use the information presented here as guidelines only. PanAmerican Seed recommends that growers conduct a trial of products under their own conditions. Crop times will vary depending on the climate, location, time of year, and greenhouse environmental conditions. It is the responsibility of the grower to confirm the treatment is available in their region as well as read and follow all the current label directions relating to the products. Nothing herein shall be deemed a warranty or guaranty by PanAmerican Seed of any products listed herein. PanAmerican Seed's terms and conditions of sale shall apply to all products listed herein.

Variety Pictures



Hot Tomato



Hot Tomato

PanAmericanSeed™

PanAmerican Seed Co.
622 Town Road, West Chicago, Illinois, USA, 60185-2698
630 231-1400 Fax: 630 231-3609 PanAmSeed.com

™ denotes a trademark of and © denotes a registered trademark of Ball Horticultural Company in the US. It may also be registered in other countries.
©2020 Ball Horticultural Company