

## Dianthus Dynasty F<sub>1</sub> Series

(*Dianthus barbatus interspecific*)

Annuals Culture (revised 01/14/21)

**Plug crop time:** 4 to 5 weeks

**Transplant to finish:** 8 to 9 weeks

- Very well-branched, upright and strong-stemmed, these excellent landscape and garden performers double as a great backyard cut flower.
- Fast-blooming plants respond well to growth regulators applied for potted plant height and tighter umbels.
- Performs as a perennial in USDA Hardiness Zones 6a to 9b (-10 to 25°F/-23 to -4°C minimum temperature).

### General Information

Exposure	Bloom Season	Height	Spread	Spacing
Sun	Early Spring, Spring, Autumn	16-20 in. (41-51 cm)	10 in. (25 cm)	10-12 in. (25-30 cm)

### Germination

Seed Form	Recommended Plug Size	Seeds/Cell	Plug Crop Weeks	Days from 50% to maximum germination	Initial Media pH/EC (1:2)	Cover Seed
PEL	288	1	5-6	3-4	5.8-6.2 pH 0.75 mmhos/cm	Yes

### Plug Production

	Stage 1	Stage 2	Stage 3	Stage 4
<b>Moisture</b>	Level 4	Level 3-4	Level 2-3	Level 2-3
<b>Temperature</b>	64-68°F (18-20°C)	65-70°F (18-21°C)	60-65°F (16-18°C)	55-60°F (13-16°C)
<b>Light</b>	Optional	5-8 mol·m <sup>-2</sup> ·d <sup>-1</sup>	8-10 mol·m <sup>-2</sup> ·d <sup>-1</sup>	10 mol·m <sup>-2</sup> ·d <sup>-1</sup>
<b>Fertiliser</b>		Less than 100 ppm N (Less than 0.7 EC)	100 to 175 ppm N (0.7 to 1.2 EC)	175 to 225 ppm N (1.2 to 1.5 EC)
<b>PGR</b>		paclobutrazol/5 ppm/Spray	paclobutrazol/5 ppm/Spray	

### Growing on to Finish

Growing on Temperature	Target Media pH/EC (1:2)	Fertiliser	Daylength
(day) 60-72°F (16-22°C) (night) 50-60°F (10-16°C)	5.8-6.2 pH 1.5-2.0 mmhos/cm	100 to 175 ppm N (0.7 to 1.2 EC)	Facultative Long Day

### Crop Scheduling

Container Size	Plugs/Pot	Crop Time	Season	PGR
306 Pack/1801	1 (ppp)	8-9 (weeks)	Late Spring	paclobutrazol 15-20 ppm Spray
4"/4.5"/Quart/10 cm	1 (ppp)	8-9 (weeks)	Late Spring	paclobutrazol 15-20 ppm Spray
5"/6"/1 Gallon/15 cm	1-3 (ppp)	8-9 (weeks)	Late Spring	paclobutrazol 15-20 ppm Spray

### Fertiliser Notation

When grown in the greenhouse, begin fertilizing the crop with 150 ppm N once a week starting a week after transplant. The plants may require additional fertilization when grown outside. If necessary, treat with a foliar spray of calcium or a calcium nitrate fertilizer supplying 150 ppm N.


## Common Problems

Insects: Watch for Aphid infestations during early growth stages.

NOTE: Growers should use the information presented here as guidelines only. PanAmerican Seed recommends that growers conduct a trial of products under their own conditions. Crop times will vary depending on the climate, location, time of year, and greenhouse environmental conditions. It is the responsibility of the grower to confirm the treatment is available in their region as well as read and follow all the current label directions relating to the products. Nothing herein shall be deemed a warranty or guaranty by PanAmerican Seed of any products listed herein. PanAmerican Seed's terms and conditions of sale shall apply to all products listed herein.

## Variety Pictures


---




Orchid




Pink Magic




Purple



Red



Rose Lace



White Blush



Mixture



PanAmerican Seed Co.  
622 Town Road, West Chicago, Illinois, USA, 60185  
+1 800-231-7065 [PanAmSeed.com](http://PanAmSeed.com)

™ denotes a trademark of and ® denotes a registered trademark of Ball Horticultural Company in the US. It may also be registered in other countries.  
©2025 Ball Horticultural Company