

Pepper (Ornamental) Hot Pops Series

(*Capsicum annuum*)

Potted Plants Culture (revised 12/02/24)

- Pump up your Autumn pot plant sales with an unusual fruit form and multicolours!
- Small, round peppers show multiple colours simultaneously to make a high-impact display for outdoor and indoor decorating.
- Compact, heavily branched plants with determinate growth.
- Perfect for quart and Mum Pals programs, plus Autumn mixed containers.

NOTE: Fruit is very hot on Hot Pops Purple, and medium hot on Hot Pops Yellow.

General Information

Exposure	Bloom Season	Height	Spread	Spacing
Sun	Summer, Autumn	4-5 in. (10-13 cm)	6-8 in. (15-20 cm)	5-7 in. (13-18 cm)

Germination

Seed Form	Recommended Plug Size	Seeds/Cell	Plug Crop Weeks	Days from 50% to maximum germination	Initial Media pH/EC (1:2)	Cover Seed
RAW	288	1	4-5	5-7	5.8-6.2 pH 0.75 mmhos/cm	No

Plug Production

	Stage 1	Stage 2	Stage 3	Stage 4
Moisture	Level 4	Level 3-4	Level 2-4	Level 2-4
Temperature	72-76°F (22-24°C)	68-73°F (20-23°C)	68-70°F (20-21°C)	68-70°F (20-21°C)
Light	Optional	1,000-2,500 f.c. (10,800-26,900 Lux)	1,000-2,500 f.c. (10,800-26,900 Lux)	2,500-5,000 f.c. (26,900-53,800 Lux)
Fertiliser		Less than 100 ppm N (Less than 0.7 EC)	100 to 175 ppm N (0.7 to 1.2 EC)	100 to 175 ppm N (0.7 to 1.2 EC)

Propagation Key Tips

Performs best under high light and warm temperatures. If necessary, uniconazole at rate of 2.5 ppm can be applied at 2 weeks after sowing for height control. Repeat 2 weeks later if needed.

Growing on to Finish

Growing on Temperature	Target Media pH/EC (1:2)	Fertiliser	Daylength
(day) 68-80°F (20-27°C) (night) 65-70°F (18-21°C)	5.5-6.2 pH 1.2-1.5 mmhos/cm	175 to 225 ppm N (1.2 to 1.5 EC)	Day Neutral

Crop Scheduling

Container Size	Plugs/Pot	Crop Time	Season	PGR
4"/4.5"/Quart/10 cm	1 (ppp)	7-13 (weeks)	Summer	-
5"/6"/1 Gallon/15 cm	1-3 (ppp)	7-13 (weeks)	Summer	-
4"/4.5"/Quart/10 cm	1 (ppp)	7-13 (weeks)	Autumn	-
5"/6"/1 Gallon/15 cm	1-3 (ppp)	7-13 (weeks)	Autumn	-

Finishing Key Tips

Performs best under high light and warm temperatures. Add 4 to 5 weeks for Spring production crop times. Grow plants dry to minimise stretch.

NOTE: Growers should use the information presented here as guidelines only. PanAmerican Seed recommends that growers conduct a trial of products under their own conditions. Crop times will vary depending on the climate, location, time of year, and greenhouse environmental conditions. It is the responsibility of the grower to confirm the treatment is available in their region as well as read and follow all the current label directions relating to the products. Nothing herein shall be deemed a warranty or guaranty by PanAmerican Seed of any products listed herein. PanAmerican Seed's terms and conditions of sale shall apply to all products listed herein.

Variety Pictures



Purple



Yellow



PanAmerican Seed Co.
622 Town Road, West Chicago, Illinois, USA, 60185
+1 800-231-7065 PanAmSeed.com

TM denotes a trademark of and [®] denotes a registered trademark of Ball Horticultural Company in the US. It may also be registered in other countries.
©2026 Ball Horticultural Company