GrowerFacts

Pepper Peppi F1 Series

(Capsicum annuum)

Vegetables Culture (revised 11/01/23)

- These small, pointed fruits offer incredible sweetness and rich flavour.
- The fruit are often seedless and are great for snacking fresh off the plant.
- To produce seedless fruit, grow it in isolation from other (non-seedless) pepper varieties.
- Best grown in-ground with support of stakes or a cage.

General Information

| Exposure | Bloom Season | Height | Spread | Spacing |
|----------|--------------|--------------------------|-------------------------|-------------------------|
| Sun | Summer | 36-48 in. (91-122 cm) | 24-36 in. (61-91 cm) | 18-24 in. (46-61 cm) |

Germination

| Seed Form | Recommended Plug Size | Seeds/Cell | Plug Crop Weeks | Days from 50% to maximum germination | Initial Media pH/EC (1:2) | Cover Seed |
|-----------|--------------------------|------------|-----------------|--------------------------------------|------------------------------|-------------|
| RAW | 288 | 1 | 5-6 | 5-7 | 5.8-6.2 pH 0.75 mmhos/cm | Light cover |

Plug Production

| | Stage 1 | Stage 2 | Stage 3 | Stage 4 |
|-------------|---|---|---|---|
| Moisture | Level 4 | Level 3-4 | Level 2-4 | Level 2-4 |
| Temperature | 70-78°F (21-26°C) | 68-72°F (20-22°C) | 65-70°F (18-21°C) | 62-67°F (17-19°C) |
| Light | Optional | 1,000-2,500 f.c. (10,800-26,900 Lux) | 1,000-2,500 f.c. (10,800-26,900 Lux) | 2,500-5,000 f.c. (26,900-53,800 Lux) |
| Fertiliser | Less than 100 ppm N (Less than 0.7 EC) | Less than 100 ppm N (Less than 0.7 EC) | 100 to 175 ppm N (0.7 to 1.2 EC) | 100 to 175 ppm N (0.7 to 1.2 EC) |

Fertiliser Notation

Pepper is very sensitive to high salts, particularly high ammonium, during germination. Keep ammonium levels to less than 10 ppm.

Propagation Key Tips

Height can be controlled by withholding fertiliser, avoiding overwatering, growing under high light conditions and using DIF when possible. If necessary, uniconazole (Sumagic) at the rate of 2.5 to 5 ppm can be applied at 2 weeks after sowing for height control. Repeat 2 weeks later if needed.

Growing on to Finish

| Growing on Temperature | Target Media pH/EC (1:2) | Fertiliser | Daylength |
|---------------------------|-----------------------------|------------------|-----------|
| (day) 68-80°F (20-27°C) | 5.5-6.2 pH | 100 to 175 ppm N | |
| (night) 65-70°F (18-21°C) | 0.75-1.0 mmhos/cm | (0.7 to 1.2 EC) | |

Days to Green Fruit

55-60

Days to Ripe Fruit

75-80

Crop Scheduling

| Container Size | Plugs/Pot | Crop Time | Season | PGR |
|---------------------|-----------|---------------|--------|-----|
| 4"/4.5"/Quart/10 cm | 1 (ppp) | 4-6 (weeks) | Spring | - |
| Field grown | 1 (ppp) | 11-12 (weeks) | Spring | - |

Finishing Key Tips

Large habit. Performs best when grown in-ground and with support.

NOTE: Growers should use the information presented here as guidelines only. PanAmerican Seed recommends that growers conduct a trial of products under their own conditions. Crop times will vary depending on the climate, location, time of year, and greenhouse environmental conditions. It is the responsibility of the grower to confirm the treatment is available in their region as well as read and follow all the current label directions relating to the products. Nothing herein shall be deemed a warranty or guaranty by PanAmerican Seed of any products listed herein. PanAmerican Seed's terms and conditions of sale shall apply to all products listed herein.

Variety Pictures





PanAmerican Seed Co. 622 Town Road,West Chicago, Illinois, USA, 60185 +1 800-231-7065 PanAmSeed.com ™ denotes a trademark of and ® denotes a registered trademark of Ball Horticultural Company in the US. It may also be registered in other countries. ©2025 Ball Horticultural Company