

## Torenia Kauai™ Series

(*Torenia fournieri*)

Annuals Culture (revised 05/15/26)

**Plug crop time:** 5 to 6 weeks

**Transplant to finish:** 5 to 7 weeks

- Compact series offers the fullest complement of the most vivid colours.
- Easy to schedule with a tight, 7 to 9-day flowering window.
- Crop management is easier and more cost-effective with this compact torenia.
- Superior performance in extreme heat and humidity.
- Brings an exotic look to small gardens, mixed combos and colour bowls; also suited to indoor containers.
- Includes unique colours Lemon Drop and Magenta.

### General Information

Exposure	Bloom Season	Height	Spread	Spacing
Shade, Partial Sun	Late Spring, Summer	8 in. (20 cm)	8 in. (20 cm)	8 in. (20 cm)

### Germination

Seed Form	Recommended Plug Size	Seeds/Cell	Plug Crop Weeks	Days from 50% to maximum germination	Initial Media pH/EC (1:2)	Cover Seed
PEL	288	1	5-6	4-6	5.5-6.0 pH 0.75 mmhos/cm	No

### Plug Production

	Stage 1	Stage 2	Stage 3	Stage 4
<b>Moisture</b>	Level 4	Level 3	Level 3-4	Level 3-4
<b>Temperature</b>	71-76°F (22-24°C)	68-73°F (20-23°C)	68-70°F (20-21°C)	65-67°F (18-19°C)
<b>Light</b>	Light	2,500 f.c. (26,900 Lux)	2,500 f.c. (26,900 Lux)	2,500-5,000 f.c. (26,900-53,800 Lux)
<b>Fertiliser</b>		Less than 100 ppm N (Less than 0.7 EC)	100 to 175 ppm N (0.7 to 1.2 EC)	100 to 175 ppm N (0.7 to 1.2 EC)

### Growing on to Finish

Growing on Temperature	Target Media pH/EC (1:2)	Fertiliser	Daylength
(day) 65-70°F (18-21°C) (night) 62-64°F (17-18°C)	5.8-6.2 pH 1.5-2.0 mmhos/cm	175 to 225 ppm N (1.2 to 1.5 EC)	Day Neutral

### Crop Scheduling

Container Size	Plugs/Pot	Crop Time	Season	PGR
Cell Pack	1 (ppp)	5-6 (weeks)	Spring	chlormequat chloride 500-700 ppm Spray
4"/4.5"/Quart/10 cm	1 (ppp)	6-7 (weeks)	Spring	chlormequat chloride 500-750 ppm Spray

### Finishing Key Tips

No pinch needed. Growth Regulators: Chlormequat chloride can be used at rate of 500 to 750 ppm (4.2 to 6.4 ml/l 11.8% formulation or 0.7 to 1.0 g/l of 75% formulation) at two weeks after transplant; repeat as necessary. Paclobutrazol 20 to 30 ppm (5.0 to 7.5 ml/l, 0.4% formulation) spray also works but is slightly less effective than chlormequat chloride. Avoid using daminozide or tank mix of daminozide/chlormequat chloride as daminozide will bleach flower colour to become less intense. Daminozide will also delay flower timing.


NOTE: Growers should use the information presented here as guidelines only. PanAmerican Seed recommends that growers conduct a trial of products under their own conditions. Crop times will vary depending on the climate, location, time of year, and greenhouse environmental conditions. It is the responsibility of the grower to confirm the treatment is available in their region as well as read and follow all the current label directions relating to the products. Nothing herein shall be deemed a warranty or guaranty by PanAmerican Seed of any products listed herein. PanAmerican Seed's terms and conditions of sale shall apply to all products listed herein.

## Variety Pictures


---



Blue & White



Burgundy




Deep Blue




Lemon Drop




Magenta



Rose



White



Mixture



PanAmerican Seed Co.  
622 Town Road, West Chicago, Illinois, USA, 60185  
+1 800-231-7065 [PanAmSeed.com](http://PanAmSeed.com)

™ denotes a trademark of and © denotes a registered trademark of Ball Horticultural Company in the US. It may also be registered in other countries.  
©2026 Ball Horticultural Company