

Euphorbia Glamour F₁

(*Euphorbia graminea*)

Annuals Culture (revised 10/31/23)

Plug crop time: 3 to 4 weeks

Transplant to finish: 5 to 8 weeks

- The best variety for in-ground performance.
- Early to flower, with the vigour to hold up to a range of weather challenges.
- An exceptional choice for a 'Baby's Breath Gypsophila' like application in the landscape.
- Use with vigorous plants like Tidal Wave® Red Velour Petunia for a stunning combination in-ground, or weave like a cloud-like hedge through more bold-textured annuals for great contrast.

General Information

Exposure	Bloom Season	Height	Spread	Spacing
Sun	Spring, Late Spring, Summer, Late Summer, Autumn, Winter	15-24 in. (38-61 cm)	15-24 in. (38-61 cm)	14-16 in. (36-41 cm)

Germination

Seed Form	Recommended Plug Size	Seeds/Cell	Plug Crop Weeks	Days from 50% to maximum germination	Initial Media pH/EC (1:2)	Cover Seed
RAW	288	1	4-5	3-6	5.8-6.2 pH 0.75 mmhos/cm	Optional
	128	2	4-5			

Plug Production

	Stage 1	Stage 2	Stage 3	Stage 4
Moisture	Level 4	Level 4	Level 3-4	Level 2-3
Temperature	65-72°F (18-22°C)	65-72°F (18-22°C)	65-72°F (18-22°C)	65-72°F (18-22°C)
Light	Optional	1,000-2,500 f.c. (10,800-26,900 Lux)	1,000-2,500 f.c. (10,800-26,900 Lux)	2,500-5,000 f.c. (26,900-53,800 Lux)
Fertiliser		Less than 100 ppm N (Less than 0.7 EC)	100 to 175 ppm N (0.7 to 1.2 EC)	100 to 175 ppm N (0.7 to 1.2 EC)
PGR			daminozide/2,500 ppm/Spray	daminozide/2,500-5,000 ppm/Spray

Propagation Key Tips

Avoid temperatures below 62°F (16°C), as cooler temperatures cause foliage yellowing. Plug sizes 128 or larger should be sown at 2 seeds per cell. Apply daminozide spray for height control, with the first application at true leaf stage, followed by a second application 7 days later. An alternative to daminozide is an application of paclobutrazol spray at 2.5 ppm or drench at 0.25-0.5 ppm at radicle emergence. This has been effective in controlling hypocotyl stretch. Glamour is vigorous, so this is a key tip. Follow either of these treatments with a single daminozide spray of 2,500 to 5,000 ppm in Stage 3.

Growing on to Finish

Growing on Temperature	Target Media pH/EC (1:2)	Fertiliser	Daylength
(day) 65-77°F (18-25°C) (night) 65-68°F (18-20°C)	5.8-6.2 pH 1.5-2.0 mmhos/cm	175 to 225 ppm N (1.2 to 1.5 EC)	Day Neutral

Crop Scheduling

Container Size	Plugs/Pot	Crop Time	Season	PGR
5'/6"/1 Gallon/15 cm	1-2 (ppp)	5-8 (weeks)	Spring	daminozide 2,500-5,000 ppm Spray
8'/2 Gallon/20 cm	2 (ppp)	6-8 (weeks)	Spring	daminozide 2,500-5,000 ppm Spray

Finishing Key Tips

Do not grow plants at temperatures below 62°F (16°C). Stretched plugs can be planted deep for better habit management. Pinching is not necessary. Daminozide may delay full flowering, but it is helpful early on in developing good branching. If starting with a daminozide spray for habit management, consider a paclobutrazol spray (10 ppm) or a low rate drench (1-2 ppm) to finish. Warmer temperatures and higher light levels can significantly reduce crop times.

NOTE: Growers should use the information presented here as guidelines only. PanAmerican Seed recommends that growers conduct a trial of products under their own conditions. Crop times will vary depending on the climate, location, time of year, and greenhouse environmental conditions. It is the responsibility of the grower to confirm the treatment is available in their region as well as read and follow all the current label directions relating to the products. Nothing herein shall be deemed a warranty or guaranty by PanAmerican Seed of any products listed herein. PanAmerican Seed's terms and conditions of sale shall apply to all products listed herein.

Variety Pictures



Glamour



Glamour



Glamour



Glamour



PanAmerican Seed Co.
622 Town Road, West Chicago, Illinois, USA, 60185
+1 800-231-7065 PanAmSeed.com

™ denotes a trademark of and © denotes a registered trademark of Ball Horticultural Company in the US. It may also be registered in other countries.
©2024 Ball Horticultural Company