

Pansy (Spreading) Cool Wave® F₁ Series

Annuals Culture (revised 10/26/23)

(*Viola x wittrockiana*)

Twice the spread...double the colour!

Wow 'em with Cool Wave – hands down, it's the *fastest-growing, longest-lasting, truly trailing pansy™* for Spring and Autumn. One trip to the garden centre for lush baskets or pots for in-ground planting is enough to convince anyone that Cool Wave is a cool-season gardener's dream.

Plug crop time: 4 to 5 weeks

Transplant to finish: Spring 6 to 7 weeks, Autumn 4 to 5 weeks

- The best-performing spreading pansy series delivers *easy spreading colour™* for Spring and Autumn, so you can extend your growing and selling season for the popular Wave® brand.
- More vigorous with the most prostrate habit of any spreading pansy – great in combos and as groundcover.
- Superior overwintering hardiness – Cool Wave is the earliest pansy to return in the Spring.
- Ideal for 4.5 to 6-in. (11 to 15-cm) pots, baskets and patio planters.

General Information

| Exposure | Bloom Season | Height | Spread | Spacing |
|----------|--|-----------------------|-------------------------|-------------------------|
| Sun | Early Spring, Spring, Late Spring, Summer, Late Summer, Autumn, Winter | 6-8 in. (15-20 cm) | 24-30 in. (61-76 cm) | 10-12 in. (25-30 cm) |

Germination

| Seed Form | Recommended Plug Size | Seeds/Cell | Plug Crop Weeks | Days from 50% to maximum germination | Initial Media pH/EC (1:2) | Cover Seed |
|-----------|-----------------------|------------|-----------------|--------------------------------------|-----------------------------|------------|
| PRM | 288 | 1 | 4-5 | 2-3 | 5.5-5.8 pH 0.75 mmhos/cm | Yes |

Plug Production

| | Stage 1 | Stage 2 | Stage 3 | Stage 4 |
|--------------------|----------------------|---|--|--|
| Moisture | Level 4 | Level 3-4 | Level 3-4 | Level 2-4 |
| Temperature | 65-70°F (18-21°C) | 65-70°F (18-21°C) | 65-70°F (18-21°C) | 60-65°F (16-18°C) |
| Light | Optional | 2,500 f.c. (26,900 Lux) | 2,500 f.c. (26,900 Lux) | 5,000 f.c. (53,800 Lux) |
| Fertiliser | | Less than 100 ppm N (Less than 0.7 EC) | 100 to 175 ppm N (0.7 to 1.2 EC) | 100 to 175 ppm N (0.7 to 1.2 EC) |
| PGR | | | daminozide/1,500-2,500 ppm/Spray daminozide/chlormequat chloride tank mix/1,500/250-2,500/500 ppm/Spray | daminozide/1,500-2,500 ppm/Spray daminozide/chlormequat chloride tank mix/1,500/250-2,500/500 ppm/Spray |

Propagation Key Tips

Adjust PGR rates and frequency of application depending on local conditions in Stages 3 and 4. Cool Wave varieties do not respond uniformly to ancymidol in plug production, so its use is not recommended. Using 105/128 plugs promotes stronger lateral growth and quicker finish. If using a 288, transplant a younger, actively growing plug that is not rootbound.

Growing on to Finish

| Growing on Temperature | Target Media pH/EC (1:2) | Fertiliser | Daylength |
|--|--------------------------------|-------------------------------------|----------------------|
| (day) 62-70°F (17-21°C) (night) 55-60°F (13-16°C) | 5.5-5.8 pH 1.2-1.5 mmhos/cm | 175 to 225 ppm N (1.2 to 1.5 EC) | Facultative Long Day |

Crop Scheduling

| Container Size | Plugs/Pot | Crop Time | Season | PGR |
|------------------------------|-----------|--------------|--------------|--|
| 5"/6"/1 Gallon/15 cm | 1-3 (ppp) | 6-7 (weeks) | Early Spring | daminozide/chlormequat chloride tank mix 2,500/250 ppm Spray |
| 10" Pot or HB/3 Gallon/25 cm | 4 (ppp) | 9-10 (weeks) | Early Spring | daminozide/chlormequat chloride tank mix 2,500/250 ppm Spray |
| 5"/6"/1 Gallon/15 cm | 1-3 (ppp) | 5-6 (weeks) | Autumn | daminozide/chlormequat chloride tank mix 5,000/500 ppm Spray |
| 10" Pot or HB/3 Gallon/25 cm | 4 (ppp) | 7-8 (weeks) | Autumn | daminozide/chlormequat chloride tank mix 5,000/500 ppm Spray |

Common Problems


Insects: Fungus Gnats and Shore Flies in plug stage and Aphids in early stages after transplant. Disease: Damping off, Black Root Rot, Foliar Leaf Spots, Botrytis

Finishing Key Tips

In heat of Autumn production, a paclobutrazol drench of 0.1 to 0.125 can be used once foliage is covering soil. Consult Cool Wave Production Handbook for more detailed information on scheduling for Autumn and Spring programs and variety-specific PGR information.

NOTE: Growers should use the information presented here as guidelines only. PanAmerican Seed recommends that growers conduct a trial of products under their own conditions. Crop times will vary depending on the climate, location, time of year, and greenhouse environmental conditions. It is the responsibility of the grower to confirm the treatment is available in their region as well as read and follow all the current label directions relating to the products. Nothing herein shall be deemed a warranty or guaranty by PanAmerican Seed of any products listed herein. PanAmerican Seed's terms and conditions of sale shall apply to all products listed herein.

Variety Pictures



Blue




Blue Skies




Fire



Frost



Golden Yellow



Lemon




Morpho




Purple




Raspberry



Raspberry Swirl



Red Wing




Strawberry Swirl




Sunshine 'N Wine



Violet Wing



White



Berries 'N Cream Mixture



Mixture



Pastel Mixture



PanAmerican Seed Co.
622 Town Road, West Chicago, Illinois, USA, 60185
+1 800-231-7065 PanAmSeed.com

™ denotes a trademark of and © denotes a registered trademark of Ball Horticultural Company in the US. It may also be registered in other countries.
©2024 Ball Horticultural Company