

Cucumber Martini F₁

(*Cucumis sativus*)

Vegetables Culture (revised 11/01/23)

- Features non-yellowing, white exterior and earlier yields.
- Very tender, "no peel" skin if harvested at 5 to 6 in./13 to 15 cm length, with white, bitter-free interior that is very sweet and crunchy.
- Best in-ground.

General Information

| Exposure | Bloom Season | Height | Spread | Spacing |
|----------|--------------|------------------------|---------------------------|-------------------------|
| Sun | - | 6-12 in. (15-30 cm) | 48-72 in. (122-183 cm) | 18-24 in. (46-61 cm) |

Germination

| Seed Form | Recommended Plug Size | Seeds/Cell | Plug Crop Weeks | Days from 50% to maximum germination | Initial Media pH/EC (1:2) | Cover Seed |
|-----------|-----------------------|------------|-----------------|--------------------------------------|-----------------------------|------------|
| RAW | Direct sow 128 | 1-2 1 | 3-4 1-2 | 2-3 | 5.8-6.2 pH 0.75 mmhos/cm | Yes |

Plug Production

| | Stage 1 | Stage 2 | Stage 3 | Stage 4 |
|--------------------|---|---|---|---|
| Moisture | Level 4 | Level 3-4 | Level 2-3 | Level 2-3 |
| Temperature | 70-75°F (21-24°C) | 65-70°F (18-21°C) | 65-70°F (18-21°C) | 65-70°F (18-21°C) |
| Light | Optional | 1,000-2,500 f.c. (10,800-26,900 Lux) | 1,000-2,500 f.c. (10,800-26,900 Lux) | 2,500-5,000 f.c. (26,900-53,800 Lux) |
| Fertiliser | Less than 100 ppm N (Less than 0.7 EC) | Less than 100 ppm N (Less than 0.7 EC) | Less than 100 ppm N (Less than 0.7 EC) | Less than 100 ppm N (Less than 0.7 EC) |

Propagation Key Tips

Best performance when directly sown into final container. Height can be controlled by withholding fertiliser, avoiding overwatering, growing under high light conditions and using DIF when possible.

Growing on to Finish

| Growing on Temperature | Target Media pH/EC (1:2) | Fertiliser | Daylength |
|--|----------------------------|-------------------------------------|-----------|
| (day) 65-70°F (18-21°C) (night) 60-65°F (16-18°C) | 5.8-6.2 pH 1.0 mmhos/cm | 100 to 175 ppm N (0.7 to 1.2 EC) | - |

Days to Ripe Fruit

55-60 (direct sow)

Crop Scheduling

| Container Size | Plugs/Pot | Crop Time | Season | PGR |
|---------------------|-----------|-------------|--------|-----|
| Cell Pack | 1-2 (ppp) | 2-3 (weeks) | Spring | - |
| 4"/4.5"/Quart/10 cm | 1-2 (ppp) | 2-3 (weeks) | Spring | - |

Finishing Key Tips

Direct-sow into final container. Performs best when grown in-ground. Vining plants can be trained up a trellis to save garden space.

NOTE: Growers should use the information presented here as guidelines only. PanAmerican Seed recommends that growers conduct a trial of products under their own conditions. Crop times will vary depending on the climate, location, time of year, and greenhouse environmental conditions. It is the responsibility of the grower to confirm the treatment is available in their region as well as read and follow all the current label directions relating to the products. Nothing herein shall be deemed a warranty or guaranty by PanAmerican Seed of any products listed herein. PanAmerican Seed's terms and conditions of sale shall apply to all products listed herein.

Variety Pictures



Martini



Martini



PanAmerican Seed Co.
622 Town Road, West Chicago, Illinois, USA, 60185
+1 800-231-7065 PanAmSeed.com

™ denotes a trademark of and © denotes a registered trademark of Ball Horticultural Company in the US. It may also be registered in other countries.
©2024 Ball Horticultural Company